



ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

CODEN: IJRSFP (USA)

International Journal of Recent Scientific Research
Vol. 9, Issue, 3(B), pp. 24745-24749, March, 2018

**International Journal of
Recent Scientific
Research**

DOI: 10.24327/IJRSR

Research Article

FUNCTIONAL RESULTS FOR ULTRA-LOW RESECTIONS OF THE RECTUM

Giorgio Maria Paolo Graziano¹, Antonio Di Cataldo² and Antonino Graziano³

¹Researcher University of Catania, Italy, Dpt Sciences Medical and Surgery Technologies Advanted

^{2,3}University of Catania, Italy Medical School, Dpt Sciences Medical Surgery and Technologies Advanted

DOI: <http://dx.doi.org/10.24327/ijrsr.2018.0903.1719>

ARTICLE INFO

Article History:

Received 05th December, 2017

Received in revised form 08th

January, 2018

Accepted 10th February, 2018

Published online 28st March, 2018

Key Words:

Incontinence new rectum

ABSTRACT

Introduction The understanding of the mechanisms concerning fecal continence has undergone a considerable impulse due to the need to reduce demolition interventions with the possibility of a definitive colostomy. (1,2,3,4) The rectus responsible for the accumulation of faecal material up to moment of defecation does not need a rectal peristaltic activity. The aim of the present study is to evaluate and analyze the functional modifications of ano rectal continence after restoration of intestinal continuity by packaging an anastomosis on the rectal abutment, or in anal colo or on the combed line. **Materials and methods:** From January 2010 to December 2017 consulted the database of the AOU polyclinic University of Catania were observed in 25 cases of neoplasia in the rectum and in 6 cases of colon neoplasia with seat: sigma (6.2%) studied at the Department of Medical Surgical Sciences and specialist II Patients selected for this analysis had a mean age of 72 years (range 74-70). All patients performed digital exploration of the rectum. The colonoscopy (fig 1.) the diagnostic imaging (CT and MRI) with which it obtained the staging of the neoplasms if. The TNM, subsequently the therapeutic path was completed by implementing the manometry. **Results:** The removal of cancer of the rectum is associated with the removal of the middle rectum and its pedicle until the origin of the inferior mesenteric artery exposes to the risk of nerve lesions with consequences on the sensibility of the residual rectum, on the sphincter function but also on the function bladder. The instrumental examinations were conducted after 6- 15-36 months from the surgical treatment. In the group of patients undergoing a curative intervention that included rectal colic anterior resections with mechanical or manual anastomosis in 16 cases (64%) and in colon anal interventions in 4 cases (13%), I noticed a variable reduction in basal sphincter tone , of the new rectum compliance only in the first 6 months post-intervention. **Discussion:** Anterior resection of the rectum involves a variety of specific symptoms such as increased frequency of evacuations, a different mode of defecation, urgency, difficulty with evacuation and continence disorders. The main causes are the decrease in the capacity and the compliance of the rectum, an alteration of the tone of the internal anal sphincter and the loss of the injured anal rectal reflex. The factors that influenced the lifestyle were: the excessive frequency, fragmentation and the incontinence. The incidence of functional disorders reached 90% in the first 6 months of the postoperative period to improve within 1-2 years. **Conclusion:** The saving of sympathetic and parasympathetic nervous structures represents the measure for the best conservation of the urinary and faecal containment and expulsive function. Attention to at least partial preservation of the internal sphincter and total removal of the rectum associated with current reconstruction techniques contribute to the reduction of the evacuation disorders of the sphincter function before and after the operation.

Copyright © Giorgio Maria Paolo Graziano *et al*, 2018, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

The understanding of the mechanisms concerning fecal continence has undergone a considerable impulse due to the need to reduce demolition interventions with the possibility of a definitive colostomy. (1,2,3,4) The rectus responsible for the accumulation of faecal material up to the moment defecation does not require rectal peristaltic activity. The use of a reservoir can be attributed to the capacity of the rectum of increasing fecal volumes without proportionally increasing the anal rectal pressure. Due to the muscular component to accommodate new volumes of relaxation. (5,6,7,8) The ability to transmit sensory information of rectal replenishment does not depend on the presence of the rectum but from the snow

terminations present in the anus muscle elevator such specific receptors and nerve fibers transport the sensation of rectal distension following parasympathetic , the branches until the confluence of the sacral plexus. the surgical treatment of the rectum of the under peritoneal rectum has imposed the affirmation of therapeutic protocols that provide for the preservation of the sphincter function, this has led to the need for control over time of the functional results in terms of anal rectal continence (9,10, it is of fundamental importance to diagnose any misunderstood functional deficits that become manifest after the surgical treatment up to invalidate the functional results of the sphincter conservative surgery. (11,12,13) The interventions that involve the total or total sub rectal resection modify the rectal compliance that decreases as

*Corresponding author: Giorgio Maria Paolo Graziano

Researcher University of Catania Italy Dpt Sciences Medical and Surgery Technologies Advanted

low and the anastomosis results in an increase in daily defecations. The aim of the present study is to evaluate and evaluate the functional modifications of ano rectal continence after restoration of intestinal continuity by packaging an anastomosis on the rectal abutment, or in anal colon or on the combed line

MATERIALS AND METHODS

From January 2010 to December 2017 consulted the database of the AOU Polyclinic University of Catania were observed in 25 cases of neoplasms to site in the rectum in 6 cases of colon neoplasms with seat: sigma (6.2%) studied at the Department of Medical and Surgical Sciences II The patients selected for this analysis had a mean age of 72 years (range 74-70) The clinical signs that s in patients observed at the clinical examination were: blood in the stool accompanied by diarrhea and constipation , asthenia, rapid weight loss and anemia. All patients performed digital exploration of the rectum. The colonoscopy. The diagnostic imaging (CT and MRI) with which it obtained the staging of the neoplasms if. The TNM, subsequently the therapeutic path was completed by implementing the manometry (fig 1,2)

the middle rectum and its peduncle until the origin of the inferior mesenteric artery exposes the risk of nerve lesions with consequences on the sensibility of the residual rectum, on the sphincter function but also on the bladder function. To evaluate the functional results of the evacuations after ultra-low resections we have adopted the classification of Parks 1 normal, 2 difficulty in the control of gas and diarrhea 3 loss of liquid stool 4 loss of normal stools. The analysis of the results was carried out in the preliminary phase with the compilation of a questionnaire in which it was possible to obtain information on the functional results with a report of the evacuative frequency. The electromanometric controls with the manometric evaluation were performed with a pneumo-hydraulic perfusion manometer in which the basal and contractile sphincter activity and their sensitivity were evaluated. In all tests the basal sphincter tone was altered with an average value of 37.2 mmHg (reference values 40-80 mmHg), symmetrical on all channels. HPZ (High Pressure Zone) length of 1.4 cm (norm 2-4.5cm) with average pressure values of 31.6 mmHg. (norm 40-120 mmHg). Voluntary contraction decreased in average increments of 47.4 mmHg (reference values 105-261 mmHg). RIAR present.




Fig 1 electrodes




Fig 2 Infusion pump



Fig 3 Traced EMG




Fig 4 contraction activity

which evaluated through catheters to continuous perfusion of morphology varies with balloons of different volume the functions of the internal anal sphincter, external and the anal rectus reflex before and after surgery. Electromyography (fig. 3,4) that studied neuromuscular dysfunctions of nervous or myogenic origin by measuring the resting potentials and during the contractile activity of the muscles in order to check the degree of validity of the sphincter activity in the preoperative, highlighting the presence of nerve lesions and in postoperative eventual iatrogenic damage on the sphincter complex and nerve fibers, through the insertion of bipolar, needle-shaped concentric electrodes inserted in the external anal sphincter, or in the rectal muscle. process diagnostic. Through opaque substances they allowed to evaluate the anal rectum angle in basal conditions and during the evacuation. . The conservative surgery operations of the sphincters performed were the RA anterior or mechanical or manual r resection, the colon anal anastomosis, the by anal, and the pull -through (bacon black) and finally the sphincter trans. In all patients the total or sub-total resection of the rectum was performed.

RESULTS

The removal of rectal cancer is associated with the removal of

Anal rectum compliance: F.S. (first sensation) normal at 50cc (v. <40cc), constant stimulation sensation (CS) at 80 cc (see 50-60cc) and Maximum Volume Tolerated (MTV) at 120cc (v.v 200-600 cc). The instrumental examinations were conducted after 6- 15-36 months from the surgical treatment. In the group of patients undergoing a curative intervention that included rectal colic anterior resections with mechanical or manual anastomosis in 16 cases (64%) and in colon anal interventions in 4 cases (13%), I noticed a variable reduction in basal sphincter tone , of the new rectum compliance only in the first 6 months post-intervention. the reduced tonic sphincter activity and the absence of the vascular-rectum inhibitory reflex did not allow to retain small faecal volumes both from the reservoir and from the intestinal lumen, since the latter were free to transit without being sensed by either the at receptors or the transactional nerve endings discrimination of the content this favored the soiling both during the waking state and especially during sleep. The complete functional recovery was obtained in 70% of cases treated within 15 -36 months of the operation. Sphincter functional recovery was slower (36 months) in patients whose anastomosis packaging was 2 cm from the anal fissure. In the remaining 30% there was good continence but the defecatory frequency exceeded three times a day. after 36 months it was possible to observe, with functional

investigations, the recovery of the internal sphincter, the adaptation of the external sphincter and the improvement of the anal-rectum sensitivity, as the rectal angle was conserved within the maximum limits. 10% of this group was incontinent. In these cases we have found the presence of iatrogenic lesions due to both the technical difficulties of execution or in the enlarged exeresis conducted for the purpose of oncological radicality (radical lymphadenectomy). Incontinence phenomena with increasing gravity were highlighted with the lack of control for gas or stool. Liquid, but never for solid stools. In patients with anal anastomosis and colonic reservoir (5cm from the anal rhyme) in 6.5% the mean defecation rate was 2 times a day and spontaneously began, in the remaining group of patients (7-9 cm from the anal rhyme) 6, 5% the urge to evacuate was caused every 2 days by an anemia. In patients who performed an anastomosis without reservoir (n 23 cases equal to 76.5%) the analysis of the results was carried out in the preliminary phase with the compilation of the questionnaire with which it was possible to evaluate the functional results of the evacuative frequency in average up to 2 times a day. After 6 months we performed the electromanometric checks. The manometric evaluation showed again an evident sphincter hypotonia the traces performed between 16.36 months baseline sphincter activity and the contractile activity normalize to detection. With the pull-through techniques (bacon, black) n 3 cases equal to 9%. We obtained positive responses in relation to the extension of the exeresis. Anal sensitivity depended on the integrity of the transactional mucosa and in the exeresis with pull-through technique there was a sensory deficit with the impossibility of discriminatory recognition of the solid, liquid rectal content. Gaseous, which was directly proportional to the extension of the mucosectomy so that the recovery of sensitivity was faster in patients with less extension.

DISCUSSION

The anterior resection of the rectum involves a variety of specific symptoms such as increased frequency of evacuations, a different mode of defecation, urgency, difficulty in evacuation and continence disorders. The main causes are the decrease in the capacity and the compliance of the rectum, an alteration of the tone of the internal anal sphincter and the loss of the injured anal rectal reflex. (13 14) The factors that influenced the lifestyle were: the excessive frequency, fragmentation and the incontinence. The incidence of functional disorders reached 90% in the first 6 months of the postoperative period to improve within 1-2 years with a decrease in the capacity of the new rectum and the alteration of sensitivity. (15 16) An incorrect mode of defecation that was reported in the questionnaire in 54%, urgency in 60% and an obstructive defecation in 10% after resection of the rectum. (17,18) the sensation of fullness of the rectum was present on average one year from exeresis in all patients undergoing anterior resection. the manometric evaluation in the previous mechanical or manual resections showed that the anal endo fall was due to direct iatrogenic damage on the anal sphincter, therefore the most reliable hypothesis of the origin of the sphincter damage was the probable maneuver of rectal eversion performed to facilitate the execution of colon anal anastomosis, resulting in a lesion of a nerve fiber (n pudendo) or direct damage to the muscle fibers (19.20) However, nerve lesions do not always occur in a complete section of the nerve fiber, but a

simple excessive traction can lead to a functional deficit (neurological praxia) even if transitory, with restitutio ad integrum in a period of time ranging from a few weeks to a few months, as in our observation. (21,22) The suture was also questioned when the latter being close to the combed line most likely to cause an internal sphincter lesion (23,24). Often a modification of the pelvic floor to which a modification of the ano-rectal angle was associated.(26,27) the surgical intervention, although generally does not modify it, makes potential incontinence manifest, especially when this perineal prolapse is associated with the deficit of the external anal sphincter. (28,29,30) For which it is useful to highlight the prolapse of the perineum underneath plan in the preoperative. (31,32) Under the anatomical-functional profile the nerve lesions can be distinguished in: 1 parasympathetic lesions; 2-sympathetic lesions; pudendal nerve injuries. The similar sympathetic nerve runs along the erect nerves that through the sacral foramina reach the pelvis and the inferior hypogastric plexus located at the front and lateral to the rectum, immediately above the endopelvic fascia that covers the elevating muscles of the anus. (33,34) Lesions at the lower hypogastric plexus level may occur during pelvic lymphadenectomy or during traction, which is performed on the lower portion of the rectum and result in bladder atony resulting in a retention of urine. (36,37,38) On the anal sphincter apparatus the influence is on the internal sphincter for the role of modulator of the sympathetic function. (39,40) The sympathetic thoraco-dorsal artery runs into the hypogastric plexus at the level of the aortic bifurcation immediately below the parietal peritoneum, and divides into the hypogastric nerves that run parallel to each artery, medially to the ureter. (41,42,43) A lesion of the hypogastric plexus is possible during an aortocaval lymphadenectomy, during the preparation of the inferior mesenteric artery, while the need for a high lateral dissection can compromise the hypogastric nerves. In the anal rectum the sympathetic thoraco-dorsal behaves similarly with an action of contraction on the internal anal sphincter and of relaxation on the rectal musculature. (44,45,46,47) One of his lesions has a negative influence on the basal sphincter tone and on the function of continence, but contributes to a reduction in the compliance of the residual rectal stump. Lesions of the sympathetic lead to a condition of detrusor hyper reflective for the action of sympathetic terminations on the alpha receptors at the level of the trine (contraction), and on the beta receptors at the detrusor level (release). (48,49,50,51,52) The pudendal nerve lesions are more rare, due to its course in part below the anus lift, but possible in case of low or very low anterior resection, and that may involve alterations of bladder, anal-perineal sensitivity and above all of the function of the striated sphincter.

CONCLUSIONS

The saving of sympathetic and parasympathetic nerve structures is the measure for the best conservation of the urinary and faecal containment and expulsive function. The attention to at least partial preservation of the internal sphincter and the total removal of the rectum associated with current reconstruction techniques contribute to the reduction of the evacuation disorders. The sphincter function before and after surgery was documented by the clinical anamnestic questionnaire in which the degree of continence in the faeces

and gases, the ability to discriminate the solid or liquid content. The study was integrated with the manometric examination that documented how from the six months patients had a good continence over 70% of cases treated minor disorders were highlighted by a partial and incomplete lesion of the thoracic dorsal nerve or muscle fibers of the internal syndromes with a reduction in compliance and continence in the residual rectal abutment. A transient functional deficiency of the duration between 12-24 months from treatment,

References

- Williamson JL, Nelson RL, Orsay C, Pearl RK, Abcarian H.A comparison of simultaneous longitudinal and radial recordings of anal canal pressures. *Dis Colon Rectum* 1990;33:201-206;
- Orrom WJ, Wong WD, Rothenberger DA, Jensen LL. Evaluation of an air filled microballoon and mini-transducer in the clinical practice of anorectal manometry; preliminary communication. *Dis Colon Rectum* 1990;33:594-597;
- Perry RE, Blatchford GJ, Christensen MS, Thorson AG, Attwood SEA. Manometric diagnosis of anal sphincter injures. *Am J Surg* 1990;159:112-117;
- Robert PL, Collier JA, Schoetz, DJ Jr, Veidenheimer MC. Manometric assessment of patients with obstetric injures and fecal incontinence. *Dis Colon Rectum* 1990;33:16-20;
- Johnson GP, Pemberton JH, Ness J, Samson M, Zinsmeister AR. Transducer manometry and the effect of body position on anal canal pressures. *Dis Colon Rectum* 1990;33:469-475;
- Sun WM, Read NW, Prior A, Daly JA, Cheah SK, Grundy D. Sensory and motor responses to rectal distension vary according to rate and pattern of balloon inflation. *Gastroenterology* 1990;99:1008-1015;
- Graziano G, et al (2017). The mystery of life. *International Journal of Advanced Research*, vol. 5, p. 2640-2646,, doi: <http://dx.doi.org/10.21474/IJAR01/3055>
- GRAZIANO G, et al (2017). The antibiotic is needed in clean surgery?. *International Journal of Recent Scientific Research*, vol. 8, p. 22339-22342, doi: <http://dx.doi.org/10.24327/ijrsr.2017.0812.1247>
- Graziano G, et al (2017). What the anti-reflux surgery ideal? *International Journal of Recent Scientific Research*, vol. 8, p. 15106-15110,, doi: 10.24327/IJRSR
- Graziano G, A Dicataldo (2017). Complications In Laparoscopic Cholecystectomy. *International Journal of Current Advanced Research*, vol. 6, p. 3855-3859, doi: <http://dx.doi.org/10.24327/ijcar.2017.3859.0382>
- Graziano G, e al (2017). Treatment For Skin of RENAL CISTS. *International Journal of Current Research*, vol. 9, p. 61178-61181,,: doi: doi.org/10.24941/ijcr.2017
- Graziano G, e al (2017). Which Treatment in the Zenker diverticulum. *International Journal of Recent Scientific Research*, vol. 8, p. 21612-21616, doi: <http://dx.doi.org/10.24327/ijrsr.2017.0811.1107>
- Cali RL, Blatchford GJ, Perry RE, Pitsch RM, Thorson AG, Christensen MA. Normal variation in anorectal manometry. *Dis Colon Rectum* 1992;35:1161-1164;
- Jorge JM, Wexner SD. Anorectal manometry: techniques and clinical applications. *South Med J* 1993;86(8):924-931;
- Meunier PD, Gallavardin D. Anorectal manometry: the state of the art. *Dig Dis* 1993;11(4-5):252-264;
- Sundblad M, Hallbook O, Sjodahl R. Anorectal Manometry with a microtransducer. *Eur J Surg* 1993;159(6-7):365-370;
- Braun JC, Treutner KH, Drew B, Klimaszewski M, Schumpelick V. Vectormanometry for differential diagnosis of fecal incontinence. *Dis Colon Rectum* 1994; 37:989-986;Loening-Baucke V. Anorectal manometry and biofeedback training. In "Pediatric Gastrointestinal Motility Disorders" Hyman PE, Di Lorenzo C eds Academy Professional Information Services, Inc. New York, 1994; pp 231-252;
- Graziano G (2016). Which Treatment In Cystic Tumors of The Pancreas: Conservative or Resection. *International Journal of Current Advanced Research*, vol. 5, p. 1190-1198, ISSN: 2319-6505, doi: DOI: 10.24327/IJCAR
- Graziano G, et al (2017). Congenital Anomalies of the Kidney and Urinary Tract Neoplasms and in the Elderly. *International Journal of Advanced Research*, vol. 5, p. 265-273, doi: DOI URL: <http://dx.doi.org/10.21474/IJAR01/3512>
- Graziano G, et al (2017). Lithiasis In Urinary Diversions or Post Prostatectomy. *International Journal of Recent Scientific Research*, vol. 8, p. 16357-16363, doi: <http://dx.doi.org/10.24327/ijrsr.2017.0804.0136>
- Graziano G, et al (2017). Papillary Bladder Tumor. *International Journal of Recent Scientific Research*, vol. 8, p. 18485-18490, doi: <http://dx.doi.org/10.24327/ijrsr.2017.0807.0518> McHugh SM, Diamant NE. Anal canal pressure profile: a reappraisal as determined by rapid pull-through technique. *Gut* 1987;28:1234-1241;
- Graziano G, et al (2016). Renal Uteroscopy Treatment of Kidney and Bladder Stones. *International Journal of New Technology And Research*, vol. 2, p. 135-138
- Graziano G, et al (2016). Vascular Thoracic Fibrous Adipose Tissue (New Disease). *Journal of Pharmaceutical And Biomedical Sciences*, vol. 6, p. 419-424, , doi: <http://dx.doi.org/10.20936/jpbms/160265>
- Graziano G, et al (2016). Clinical and Molecular Anatomy of Gastrointestinal Stromal Tumors (GIST) *International Journal of New Technology And Research*, vol. 2, p. 110-114
- Graziano G, et al (2016). Early Epithelial Ovarian Carcinoma Treatment (IF 2.995). *International Journal of New Technology and Research*, vol. 2, p. 69-74, ISSN: 2454-411615.
- Graziano G, et al (2016). On Traumatic Lesions of The Pancreas (IF 2.09). *World Journal of Research And Review*, vol. 2, p. 24-28, ISSN: 2455-3956
- Hemond M, Bedard G, Bouchard H, Arhan P, Waitier A, Devroede G. Step-by-step anorectal manometry: small balloon tube. In "Practical guide to anorectal testing. End Edition" Smith LE, Ed Igaku-Shoin medical publishers, Inc. – New York 1995; pp101-141;
- Stein BL, Roberts PL. Manometry and the rectoanal inhibitory reflex. In: "Constipation, Etiology, Evaluation

- & Management” Wexner SD, Bartolo DCC eds. Butterworth-Heinemann – Oxford, 1995; pp 63-76;
29. Collier JA, Sangwan YP. Computerized anal sphincter manometry performance and analysis. In: “practical guide to anorectal testing 2nd edition” Smith Le ed. Igaku-Shoin Medical publishers, Inc. – New York 1995; pp 51-100;
30. Yoshiota K, Keighley MRB. The position of the patient does not adversely influence the results of the most clinically important measurements of anorectal function. *Int J Colorect Dis* 1995;10:47-48;
31. Diamant NE, Kamm MA, Wald A, Witehead WE. AGA technical review on anorectal testing techniques. *Gastroenterol* 1999; 116:735-77.
32. Collier JA. Clinical application of anorectal manometry. *Gastroenterol Clin North Am* 1987;16(1):17-33;
33. Graziano G, et al (2016). Treatment Therapies in Renal Cell Carcinoma in elderly: A Descriptive Analysis (IF 2.385). *International Multispecialty Journal of Health*, vol. 2, p. 20-24, ISSN: 2395-6291, doi: DOI:10.25125/medical-journal
34. Graziano G, et al (2015). The Neuroendocrine Cancer. Personal Comments and Operational Remarks. *Journal of Surgery And Surgical Research*, vol. 1, p. 53-58, doi: DOI: 10.17352/2455-2968.00001418.
35. Graziano G, et al (IF 3.75) (2016). The Familial Adenomatous Polyposis. A Difficult Problem, Between Prevention and Treatment. *Journal Of Surgery And Surgical Research*, vol. 2, p. 05-09, doi: doi:10.17352/2455-2968-000021
36. Graziano G, et al (2017). Role of Genetic Mutations in the Diagnosis of Gallbladder neoplasms. *International Journal of Recent Scientific Research*, vol. 8, p. 20908-20913, DOI: <http://dx.doi.org/10.24327/ijrsr.2017.0810.098220>.
37. Graziano G, et al (2017). Single Accessed Gallbladder Surgery. *International Journal of Recent Scientific Research*, vol. 8, p. 19359-19362, DOI: <http://dx.doi.org/10.24327/ijrsr.2017.0808.0679>
38. Graziano G, et al (2017). The Use Of Bar In Colorectal Surgery In The Elderly. *International Journal of Recent Scientific Research*, vol. 8, p. 19950-19954, DOI: <http://dx.doi.org/10.24327/ijrsr.2017.0809.0793>
39. Graziano G, et al (2016). One Time Surgery in Contemporary Diseases of the Abdominal Wall and Pelvis in the Elderly. *Journal of Surgery and Surgical Research*, vol. 2, p. 18-20, doi: 10.17.352/2455-2968-00002456.
40. Graziano G, et al (2015). The Stent Evolution in Colo-Rectal Emergencies. *Journal of Surgery and Surgical Research*, vol. 1, p. 45-48, doi: 10.17352/2455-2968-00001224.
41. Graziano G, et al (IF 3, 35) (2016). Which Surgery in Geriatric Breast Cancer. *Journal of Surgery And Surgical Research*, vol. 2, p. 014-017, doi: 10.17352/2455-2968-000023
42. Giorgio Maria Paolo Graziano.2016, Diagnostic and Therapeutic In the Intestinal Duplication. *Int J Recent Sci Res.* 7(8), pp.13000-13003.
43. Graziano Giorgio Maria Paolo et al.2016, Essentiality 'In The Doctor-Patient Relationship. *Int J Recent Sci Res.* 7(12), pp.14527-14537.
44. Graziano Giorgio Maria Paolo et al A Descriptive Study of Differentially Placed Hydatid cysts *International Multispecialty Journal of Health (IMJH)* ISSN: [2395-6291] [Vol-2, Issue-4, April- 2016]
45. Graziano Giorgio maria paolo et al Which Therapeutic Treatment in Gastric Lymphoma. *World Journal of Research and Review (WJRR)* ISSN:2455-3956, Volume-2, Issue-6, June 2016 Pages 06-09
46. Giorgio Maria Paolo Graziano et al On Traumatic Lesions of The Pancreas. *World Journal of Research and Review (WJRR)* ISSN:2455-3956, Volume-2, Issue-6, June 2016 Pages 24-28
47. Marc A. Gladman, M.R.C.O.G., M.R.C.S. (Eng), S. Mark Scott, Ph.D., Christopher L.H. Chan, F.R.C.S., Norman S. Williams, M.S., F.R.C.S., Peter J. Lunniss, M.S., F.R.C.S.: “Rectal Hyposensitivity. Prevalence and Clinical Impact in Patients With Intractable Constipation and Fecal Incontinence”. D.C.R. 2003 Vol.46, N°2: 238-246-2)
48. Christopher L.H. Chan, F.R.C.S., S. Mark Scott, Ph.D., Norman S. Williams, F.R.C.S., Peter J. Lunniss, F.R.C.S. “Rectal Hypersensitivity Worsens Stool Frequency, Urgency and Lifestyle in Patients With Urge Fecal Incontinence”. D.C.R. 2005 Vol. 48, N°1: 134-140.
49. Gloria Lacima, M.D., Miguel Pera, M.D., Josep Valls-Solé, M.D., Xavier Gonzales-Argenté, M.D., Montserrat Puig-Clota, M.D.: “Electrophysiologic Studies and Clinical Findings in Females With Combined Fecal and Urinary Incontinence: A prospective Study”. D.C.R. 2006 Vol. 49 N° 3: 353-359
50. M.D. Crowell, Ph.D., B.E.Lacy, M.D., Ph.D., V.A. Schettler, B.S.N., T.N. Dineen, M.D., K.W.Olden, M.D., N.J. Talley, M.D., Ph.D.: “Subtypes of Anal Incontinence Associated With Bowel Dysfunction: Clinical, Physiologic, and Psychosocial Characterization”. D.C.R. 2004 .Vol. 47 N° 10: 1627-1635.
51. .Giorgio Maria Paolo Graziano et al.2018, The Radical Anal Trans Excision In The Initial Neoplasm of The Rectum. *Int J Recent Sci Res.* 9(2), pp. 24013-24017. DOI: <http://dx.doi.org/10.24327/ijrsr.2018.0902.1581>
52. Giorgio Maria Paolo Graziano et al. 2018, Local Recurrences After Ultra Low Resection of The Rectum. *Int J Recent Sci Res.*9(2), pp. 24119-24124. DOI: <http://dx.doi.org/10.24327/ijrsr.2018.0902.1601>.
