



**RESEARCH ARTICLE**

**INFLUENCE OF TITANIUM DIOXIDE NANOPARTICLES ON THE PHOTOSYNTHETIC AND BIOCHEMICAL PROCESSES IN ORYZA SATIVA**

**Maria Vera J. Da Costa and \*Prabhat K. Sharma**

Department of Botany, Goa University, Goa

**ARTICLE INFO**

**Article History:**

Received 14<sup>th</sup>, December, 2014

Received in revised form 23<sup>th</sup>, December, 2014

Accepted 13<sup>th</sup>, January, 2015

Published online 28<sup>th</sup>, January, 2015

**Key words:**

Ascorbic acid, Fv/Fm, Lipid peroxidation, Osmotic stress, Photosynthesis, uptake

**ABSTRACT**

The phytotoxicity of titanium dioxide nanoparticles (TiO<sub>2</sub> NPs) is a rapidly expanding area of research with a dearth of information regarding its toxicity in plants. We assessed the TiO<sub>2</sub> NP (<25 nm) effect on growth and photosynthesis in *Oryza sativa* var. Jyoti grown in a hydroponic system. Our results show that TiO<sub>2</sub> NPs do not affect germination, however, an overall decrease in seedling growth (root, shoot length and biomass) was observed at high concentration. TiO<sub>2</sub> NP exposure led to greater accumulation of Ti in roots than in shoots, decreasing CO<sub>2</sub> fixation, transpiration rate and stomatal conductance at 1000 ppm with insignificant effect on photosynthetic pigments, quantum efficiency of PSII (Fv/Fm ratio) and photochemical quenching. High treatment caused oxidative and osmotic stress evidenced by increased MDA and proline content. This study thus indicates that Ti accumulation caused phytotoxicity to growth and photosynthesis leading to oxidative and osmotic stress in rice.

© Copy Right, IJRSR, 2014, Academic Journals. All rights reserved.

**INTRODUCTION**

Metal oxide nanoparticles (NPs) are materials with size range of 1-100 nm (Klaine *et al.*, 2008). Titanium dioxide (TiO<sub>2</sub>) NPs are one of the most commonly found particles in the environment as a result of their large scale production and application for commercial purposes such as cosmetics, sunscreens (Mu *et al.*, 2010), semiconductors, pharmaceuticals, energy storage, catalysts, coatings and paints (Kaegi *et al.*, 2008), due to its UV absorption efficiency and transparency to visible light (Franklin *et al.*, 2007). It is also used in the production of textiles (Yuranova *et al.*, 2007) and for solar-driven self-cleaning coatings (Cai *et al.*, 2006). These usages would inevitably lead to increased NP entry into the environment. As far as concentration of NPs in soil or aquatic system is concerned, no direct published data is available (Navarro *et al.*, 2008). Using fate models, Mueller *et al.* (2008), estimated the concentration of TiO<sub>2</sub> NP to be 120 µg/kg<sup>3</sup> of agricultural land per year as a result of contaminated sewage sludge. Boxall *et al.* (2007) used simple box models to estimate the amount of Ti in the environment and reported that the expected amounts of Ti and other common NPs (Ce, Zn and Ag) were found to be in the range of 1 to 10 µg/L in natural waters, with total NP concentration scaling up to 100 µg/L approximately in the aquatic system currently which may increase many fold in the near future.

Release of TiO<sub>2</sub> NP in the environment is ameliorating at a very rapid rate contaminating soil and aquatic ecosystems (Farre *et al.*, 2009; Tourinho *et al.*, 2012). TiO<sub>2</sub> NPs are likely to persist and accumulate in the soil and water due to low

dissolution (Baun *et al.*, 2008) forming stable aggregates (French *et al.*, 2009). Agglomeration and aggregation, in the environment are influenced by NP surface characteristics and particle size. These are negatively correlated to soil characteristics such as dissolved organic matter, and positively correlated to pH and ionic strength (Fang *et al.*, 2009).

The mechanism of NP uptake by plant roots is not clearly understood. It is reported that depending on the size nanoparticles, NPs may enter the plant cells through carrier protein, aquaporin, ion channels, fluid phase endocytosis, plasmodesmata transport or entry facilitated through natural organic matter or root exudates and formation of new pores (Rico *et al.*, 2011). Once inside the cell, NPs can bind to the organelles or protein (Rosenbluh *et al.*, 2011) or form aggregates (Guzman *et al.*, 2006). Interaction of NPs in plants not only depends upon the surface characteristics, particle size, temperature, pH and ionic strength (Maynard *et al.*, 2006; Wiesner *et al.*, 2006) but also on the type of interactions with lipids and proteins such as electrostatic, hydrogen bonding, and hydrophobic interactions (Hug and Sulzberger, 1994; Ojamae *et al.*, 2006).

Reports on uptake of TiO<sub>2</sub> NP in plants are limited, especially in food crops (Rico *et al.*, 2011). Uptake, translocation and cell distribution of ultra small TiO<sub>2</sub> NPs (<5 nm) in *Arabidopsis thaliana* was reported to be either facilitated or inhibited by the pectin hydrogel capsule which was released by the roots (Kurepa *et al.*, 2010). Increased accumulation or inactivation of toxic heavy metals by the rhizosphere was also reported to be dependent on plant species (Watanabe *et al.*, 2008). TiO<sub>2</sub>

\*Corresponding author: **Prabhat K. Sharma**

Department of Botany, Goa University, Goa

NPs toxicity caused reduction in root cell diameter, hydraulic conductivity, transpiration and leaf growth in maize (Asli and Neumann, 2009) and produced reactive oxygen species (ROS) within organisms due to its photocatalytic activity upon UV exposure (Kahru *et al.*, 2008). TiO<sub>2</sub> NPs are reported to be beneficial to some organisms. For instance, in spinach, TiO<sub>2</sub> NPs increased chlorophyll synthesis, photosynthesis and plant dry weight (Zheng *et al.*, 2005; Hong *et al.*, 2005).

The multidisciplinary application of nanotechnology-products results in the inevitable discharge of NPs into the environment (Navarro *et al.*, 2008). Problems and challenges arise due to phytotoxicity exhibited by the NPs owing to their physicochemical surface characteristics upon interaction with the living system. The lack of knowledge and limited information on the fate and bioavailability of NPs in the environment also augment these problems (Klaine *et al.*, 2008). Our study provides a better understanding of the interaction of TiO<sub>2</sub> NP in *Oryza sativa* var. Jyoti and suggests that light reaction of photosynthesis was uninfluenced on treatment with TiO<sub>2</sub> NPs, while growth (germination, root/ shoot length and biomass) was reduced at highest concentration. We report a decline in the net photosynthesis (A, E and gs), and enhanced oxidative and osmotic stress indicated by increased MDA and proline respectively due to the TiO<sub>2</sub> NP treatment. Results also suggest generation of ascorbate in order to metabolise ROS produced, probably to mitigate the oxidative damage.

## **MATERIALS AND METHODS**

### ***Plant materials and growth conditions***

*Oryza sativa* (var. *Jyoti*), a high yielding, rice variety was obtained from Indian Council of Agricultural Research (ICAR), Goa and stored in dark at 4°C. TiO<sub>2</sub> NP (<25 nm size, Sigma-Aldrich) solution of 0, 2.5, 10, 50, 100 and 1000 ppm was prepared in Hoagland's solution (pH 6.5). Seeds were surface sterilized using 4% sodium hypochlorite (BDH) solution for 3 min, washed thoroughly with tap water and soaked for 3 days. 100 seeds were sown on germination paper in a Petri plate with respective concentration of TiO<sub>2</sub>.

The seeds were germinated in a growth chamber equipped with lamps to provide an irradiance of 200  $\mu\text{mol m}^{-2}\text{s}^{-1}$  with 16 h photoperiod, temperature of  $25 \pm 2^\circ\text{C}$  and relative humidity of 70- 75%. The 6 day old seedlings were transferred carefully in a hydroponic system containing respective concentration of TiO<sub>2</sub> NPs and were allowed to grow for a further 24 days. The hydroponic system ensured complete mixing of NPs by use of magnetic stirrers and proper aeration achieved by air pumps. The solutions were replenished every third day.

### ***Nanoparticle Characterization Studies***

TiO<sub>2</sub> NPs were obtained from Sigma-Aldrich Co. (USA). The physical characteristics as obtained from the supplier were: particle size (<25 nm), surface area (45-55  $\text{m}^2\text{g}^{-1}$ ), mp (1825 °C), and density (3.9  $\text{g mL}^{-1}$  at 25 °C), containing 99.7% trace metals basis.

### ***Size characterization using Scanning Electron Microscopy (SEM)***

TiO<sub>2</sub> NP was characterized at 30000 X magnification using SEM (JSM 5800 LV, JEOL, Japan) at the National Institute of Oceanography, Goa.

### ***Size characterization using X-ray Diffractometer (XRD)***

X-ray Diffractometer (Rigaku) pattern of TiO<sub>2</sub> NPs was recorded at room temperature in the range of  $2\theta = 20^\circ - 80^\circ$  with a step of  $0.02^\circ$  and a speed =  $2^\circ/\text{min}$  using Cu  $\alpha$  radiation ( $\lambda = 0.15406 \text{ nm}$ ).

### ***Size characterization using dynamic light scattering (DLS) analyzer***

DLS measurements in aqueous system were studied using dynamic light scattering ZetaPALS Analyzer (Brookhaven Instruments Corporation, New York, USA). Appropriate amount of TiO<sub>2</sub> NPs was suspended in distilled water as well as Hoagland's nutrient solution and was physically dispersed using an ultrasonicator for 2 min. Each measurement consisting of five replicates were measured with 10 s duration at 25°C and analyzed using BIC Particle Sizing Software Ver.5.23.

### ***Germination, root and shoot length and biomass studies***

Percent germination, root, shoot length was manually taken at the early stage (6<sup>th</sup> day) of growth. The biomass was measured taking 10 plants from each TiO<sub>2</sub> NP treatment group after 30 days of growth. Fresh leaf tissue was dried at 75°C for 72 h noted as dry weight (d.w.).

### ***Nanoparticle uptake***

Uptake of TiO<sub>2</sub> NPs was determined using an inductively coupled plasma atomic emission spectrophotometer (Perkin-Elmer Optima 2000 DV ICP-OES) calibrated with single-element calibration standards. The ash was dissolved in 1% HNO<sub>3</sub> (Merck) solution and filtered through 0.45  $\mu\text{m}$  Whatman filter paper. The filtered sample was used for ICP-OES analysis.

### ***Photosynthesis measurements***

Photosynthesis measurements were carried out using a portable infra-red gas analyzer (IRGA, ADC Bioscientific, LCi-SD, Hansatech, UK) according to Sharma and Hall (1996). The photosynthesis rate (A), transpiration rate (E) and stomatal conductance (gs) were measured at ambient temperature and carbon dioxide concentration and light intensity of  $1200 \mu\text{mol m}^{-2} \text{ s}^{-1}$  using a detachable light source provided by a dichroic lamp (Hansatech, UK).

### ***Fluorescence measurements***

Measurements were carried out using a fluorescence monitoring system (FMS, Hansatech instruments) according to Sharma *et al.* (1997) using Modfluor32 software. The maximum fluorescence (Fm), variable fluorescence (Fv) and the maximum quantum yield (Fv/Fm) ratio was calculated. The relative contributions for photochemical and non-photochemical energy dissipation measured as the photochemical quenching (qP) and non-photochemical quenching (qNP) were also calculated according to Schreiber *et al.* (1986).

### ***Extraction and analysis of photosynthetic Pigments***

Plant tissue (0.1 g) was ground in 100 % Acetone (Merck HPLC grade) making a final volume of 1.5 ml and incubated overnight at 4°C. The homogenate was centrifuged at 4°C for 10 mins at 4000 x g. 1 ml of supernatant was collected and filtered through 0.2  $\mu\text{m}$  Ultipor<sup>®</sup>N66<sup>®</sup>Nylon 6, 6 membrane

filter and 10 µl of sample was analyzed in a Waters HPLC system, according to Sharma and Hall (1996). The HPLC profile was extracted at 445 nm with a Waters 2996 photodiode array detector and peaks were identified based on RT and their spectral characteristics. -carotene was used as an external standard for peak quantization.

**Lipid per oxidation**

Lipid peroxidation of the cell membrane was determined according to Sankhalkar and Sharma (2002). Peroxidation of lipids was measured as malondialdehyde (MDA) content using an extinction coefficient of 155 mM<sup>-1</sup>cm<sup>-1</sup> at 532 nm.

**Proline determination**

Proline accumulation in fresh leaves was determined according to the method of Bates *et al.* (1973). L-Proline (Sigma ultra 99 % pure) was used as a standard for quantification by plotting a standard graph against concentration at 520 nm.

**Ascorbate assay**

Ascorbate (ASA) and dehydroascorbate (DHA) was estimated according to Kampfenkel *et al.* (1995). Ascorbic acid (sigma) was used as standard.

**Statistical analysis**

Data obtained was statistically analyzed using the procedure of ANOVA: Two-Factor without Replication and unpaired two-tailed student's t-test. Statistically significant difference 'a' is reported when the *p* < 0.05.

**RESULTS**

**Nanoparticle characterization**

The TiO<sub>2</sub> NPs were characterized under scanning electron microscope (SEM) for their shape and was observed to be spherical to round (Fig 1a). X-ray diffraction (XRD) patterns were analyzed using Scherrer's equation and the average crystalline size was measured to be 15 ± 2 nm (Fig 1b). Using dynamic light scattering (DLS) analyzer, a narrow distribution of the particle size of nano-TiO<sub>2</sub> powder in Hoagland's solution was measured to be 23 ± 2 nm (Polydispersity of 0.074) and in distilled water was in the range of 12 ± 2 nm (Polydispersity of 0.191) which were in agreement with the XRD measurements. The sizes obtained were as specified by the supplier (TiO<sub>2</sub> NPs, size <25 nm; Sigma-Aldrich, USA). The TiO<sub>2</sub> powder resulted in a suspension that appeared cloudy and did not seem to settle over the sampling time.

**Table 1** Percent germination (%G), Root and shoot length of rice plantlets (6 day old) treated with TiO<sub>2</sub> NP at different concentration.

TiO <sub>2</sub> NPs (in ppm)	% G	Root length (mm)	Shoot length (mm)
0	94.3 ± 0.9	51.31 ± 4.0	37.9 ± 1.7
2.5	93.1 ± 0.2	51.82 ± 2.6	36.4 ± 2.9
10	92.1 ± 1.4	53.04 ± 4.2	36.8 ± 2.9
50	90.9 ± 2.5	52.67 ± 1.7	36.3 ± 0.1
100	92.9 ± 2.5	47.93 ± 5.5	37.0 ± 5.3
1000	91.8 ± 1.4	39.29 ± 7.4	33.7 ± 8.7

Values are the mean (± S.D) of 7 replicates.

**Percent germination and root/ shoot length measurements**

The percentage germination was unaffected while shoot and root length decreased by approximately 11% and 23% respectively as a result of 1000 ppm TiO<sub>2</sub> NP treatment (Table 1). The shoot biomass (f.w. & d.w.) showed no change up to

100 ppm but decreased by approximately 9% at 1000 ppm (Table 2). The root biomass (f.w.) decreased slightly at higher concentration of TiO<sub>2</sub> NP (Table 2). However, in contrast to these results, the root biomass on dry weight basis increased by 476% at the highest concentration of TiO<sub>2</sub> as compared to control (Table 2). Plants grown at 1000 ppm of TiO<sub>2</sub> also showed necrosis at the leaf tips (data not shown).

**Table 2** Biomass of rice plantlets (30 day old) treated with TiO<sub>2</sub> NP at different concentration.

TiO <sub>2</sub> NPs (ppm)	Biomass (g)			
	Shoot (F.W)	Shoot (D.W)	Root (F.W)	Root (D.W)
0	0.74 ± 0.03	0.076 ± 0.02	0.33 ± 0.03	0.026 ± 0.003
2.5	0.72 ± 0.09	0.075 ± 0.02	0.32 ± 0.02	0.028 ± 0.003
10	0.76 ± 0.07	0.075 ± 0.01	0.29 ± 0.05	0.021 ± 0.003
50	0.76 ± 0.01	0.075 ± 0.01	0.30 ± 0.01	0.023 ± 0.001
100	0.77 ± 0.05	0.076 ± 0.02	0.30 ± 0.03	0.15 ± 0.02
1000	0.67 ± 0.15	0.069 ± 0.02	0.32 ± 0.13	0.13 ± 0.02

Values are the mean (± S.D) of 7 replicates.

**Accumulation of TiO<sub>2</sub> NPs using ICO-OES**

Plants showed a 3-fold increase in the Ti content up to 10 ppm as a result of the treatment (Table 3). Further increase in the TiO<sub>2</sub> concentration during treatment showed little difference in Ti content as observed in control shoots. Ti content in roots, however, showed a linear increase. 50 ppm TiO<sub>2</sub> NP treatment resulted in 3-fold increase in Ti content in roots, which was increased to 23.5 fold at 1000 ppm concentration (Table 3).

**Table 3** Shoot and root Ti content of 30 day old plantlets exposed to TiO<sub>2</sub> NP treatment.

TiO <sub>2</sub> NP (ppm)	Ti content (mg/L)	
	Shoot	Root
0	0.9 ± 0.0009	0.9 ± 0.0003
2.5	3.0 ± 0.0034	1.1 ± 0.0224
10	3.0 ± 0.0038	2.0 ± 0.0073
50	0.9 ± 0.0015	3.1 ± 0.0295
100	1.0 ± 0.0028	21.2 ± 0.914
1000	1.2 ± 0.0062	21.3 ± 0.034

Values are the mean (± S.D) of 3 replicates.

**Table 4** Carotenoid and chlorophyll content of TiO<sub>2</sub> NP treated rice plants at 0-1000 ppm concentration after 30 days of growth.

Pigments (conc. in µg)	Treatment with TiO <sub>2</sub> NP (in ppm)				
	0	2.5	10	100	1000
Neoxanthin	0.82	0.87	1.63	1.34	1.10
Violaxanthin	1.67	1.82	3.39	2.97	2.41
Antheraxanthin	0.25	0.31	0.50	0.50	0.32
Lutein	3.44	3.07	7.25	6.09	4.57
Chlorophyll b	6.72	6.77	13.54	11.87	8.93
Chlorophyll a	5.62	5.67	11.88	10.36	7.76
-Carotene	1.51	1.32	3.61	3.09	2.17

**Photosynthetic studies**

Chlorophyll fluorescence indicative of light reaction of photosynthesis was not significantly affected on treatment with high concentration of TiO<sub>2</sub> NP (Fig 2). The photosynthetic efficiency of PSII, measured as F<sub>v</sub>/F<sub>m</sub> ratio and photochemical quenching was similar to that of control, however, a 12% increase in non- photochemical quenching (qNP) indicative of dissipation of absorbed light energy in the form of heat was seen as a result of TiO<sub>2</sub> treatment (Fig 2).

The rate of CO<sub>2</sub> exchange measured as net photosynthesis (A), transpiration rate (E) and stomatal conductance (gs) decreased at higher concentrations of TiO<sub>2</sub> NP treatment (Fig 3). A slight increase in the rate of net photosynthesis (11%), transpiration rate (46%) and stomatal conductance (70%) was observed at low concentration (2.5 ppm) of TiO<sub>2</sub> NPs (Fig 3), however, an

increase in the concentration of TiO<sub>2</sub> resulted in decreased photosynthesis, transpiration and stomatal conductance. Net photosynthesis declined to 53% at 1000 ppm of TiO<sub>2</sub> concentration as compared to control. Likewise, transpiration rate and stomatal conductance also declined to 77% and 66% respectively at 1000 ppm TiO<sub>2</sub> compared to control (Fig 3).

The HPLC chromatogram obtained at 445 nm showed no qualitative changes (Fig 4). However, quantitative changes in the pigments were observed (Table 4). Chlorophyll *a* and *b* as well as carotenoid content (neoxanthin, violaxanthin, antheraxanthin, lutein, and  $\beta$ -carotene) increased on treatment with TiO<sub>2</sub> NP as compared to control (Table 4). A linear increase of up to 5% in Chl *a/b* ratio was observed as compared to control at high concentration. The Chl: Car ratio also increased at low concentration (2.5 ppm) but was not significantly different from control at high concentrations.

**Lipid peroxidation and Osmotic stress**


The malondialdehyde (MDA) content indicative of osmotic stress did not change significantly up to 50 ppm, however, an increase of 29% at 100 ppm and above concentration of TiO<sub>2</sub> NP was observed (Fig 5a).

The treatment also resulted in osmotic stress indicated by the increase in proline content (Fig 5b). Proline level increased more than 2-fold as compared to the control as a result of the TiO<sub>2</sub> treatment (Fig 6b).

Content of ascorbate, a non-enzymatic antioxidant, was also determined and was found to increase by 260% as a result of 1000 ppm TiO<sub>2</sub> treatment (Fig 5c).


**Figure Legends**

**Figure 1** Scanning electron micrograph of TiO<sub>2</sub> NP at 30000X magnification (a), X-Ray Diffraction pattern of TiO<sub>2</sub> NP (b) recorded at room temperature in the range 2 $\theta$  =20 $^{\circ}$ -80 $^{\circ}$  with a step of 0.02 $^{\circ}$ , speed= 2 $^{\circ}$ /min using Cu k radiation ( $\lambda$  =0.15406 nm).




**Figure 1** Scanning electron micrograph and XRD pattern of TiO<sub>2</sub> NP

**Figure 2** Photosynthetic efficiency of photosystem II (Fv/Fm), photochemical quenching (qP) and non- photochemical quenching (qNP) of rice plants treated with TiO<sub>2</sub> NPs after 30 days of growth. The error bars represent the mean ( $\pm$  S.D) of 7 replicates. The statistical analysis was performed using the procedure of Students t-test, 'a' representing statistically significant values when p < 0.05.




**Figure 2** Light reactions of photosynthesis

**Figure 3** Net photosynthesis rate (A), transpiration rate (E) and stomatal conductance (gs) of rice plants treated with TiO<sub>2</sub> NPs after 30 days of treatment. The error bars represent the mean ( $\pm$  S.D) of 7 replicates. The statistical analysis was performed using the procedure of Students t-test, 'a' representing statistically significant values when the p < 0.05.




**Figure 3** Dark reaction of photosynthesis

**Figure 4** HPLC profile of carotenoid and chlorophyll content of rice plants at 445 nm treated with TiO<sub>2</sub> NP after 30 days of treatment [a= Control and b= 1000 ppm TiO<sub>2</sub> NPs].



**Figure 4**HPLC chromatogram of photosynthetic pigments

**Figure 5** The MDA content (a), proline content (b) and ascorbate content (c) of rice plants treated with TiO<sub>2</sub> NP after 30 days of treatment. The error bars represent the mean ( $\pm$  S.D) of 7 replicates. The statistical analysis was performed using the procedure of Students t-test, 'a' representing statistically significant values when the p < 0.05.



**Figure 5** The MDA content (a), proline content (b) and ascorbate content (c) of rice plants treated with TiO<sub>2</sub> NP

## DISCUSSION

Our result showed no effect of TiO<sub>2</sub> NP treatment on seed germination of *Oryza sativa* var. Jyoti plants (Table 1). This may be due to the impermeability of rice seed coat to the nanoparticles till the radicle emerges. Wierzbicka and Obidzinska (1998) have also shown impermeability of seed coat to NPs. We also observed that the shoot and root length was uninfluenced up to 100 ppm, while toxicity was observed at highest concentration of TiO<sub>2</sub> NPs (Table 1). Shoot and root biomass showed no changes on fresh weight (f.w.) basis, however an increase in the root dry weight (d.w.) was observed at high concentration (Table 2). Increase in root d.w. probably may be as a result of significant accumulation of Ti in the plant roots treated with 100 and 1000 ppm TiO<sub>2</sub> NPs. Kurepa *et al.* (2010) reported that roots of *A. thaliana* released mucilage forming a hydrogel capsule either inhibiting or facilitating the entry of TiO<sub>2</sub> NPs. Watanabe (2008) reported that mucilage released may absorb, inactivate or accumulate toxic heavy metals in the rhizosphere depending on the plant species. Our data revealed massive accumulation of Ti particles in roots (Table 3), but little accumulation in shoots at high concentration. This accumulation indicates active exclusion of Ti by root cells in order to protect the shoots and more importantly the photosynthetic tissue. Ti accumulation has been reported in parenchyma of wheat root and vascular tissue (Laure *et al.*, 2011) and roots and outer leaves of cabbage plants (Hara *et al.*, 1976).

TiO<sub>2</sub> NPs did not show toxicity to light reaction of photosynthesis probably because it behaves more like metal oxides, rather than like free metal ions (Farrè *et al.*, 2009) which are known to cause toxicity. In our study, no significant changes in the PSII quantum efficiency (Fv/Fm ratio) and photochemical quenching (qP) were observed (Fig 2), however, CO<sub>2</sub> fixation rate, transpiration rate and stomatal conductance declined significantly (Fig 3) at high concentration. Our results indicate that decrease in the net CO<sub>2</sub> fixation is probably a result of reduction in the stomatal conductance rather than metabolic limitation. TiO<sub>2</sub> NP treatment results in blocking pores in addition to decreasing root pores (Asli and Neumann, 2009). This may bring about lowering of water uptake causing osmotic stress, as observed in our study (Fig 5b). This osmotic stress could lead to stomatal closure which in turn would lower the CO<sub>2</sub> fixation. In contrast to our result, Yang and Watts (2005) have reported improved growth in spinach due to TiO<sub>2</sub> treatment, in addition to, better absorption of nitrate. This then accelerates the biosynthesis of proteins and chlorophyll and enhances photosynthesis and plant biomass. Lu *et al.* (2002) also observed that TiO<sub>2</sub> NPs increased nitrate reductase activity in soyabean. We observed an increase in photosynthetic pigments as a result of TiO<sub>2</sub> but negligible increase in shoot biomass or CO<sub>2</sub> fixation.

TiO<sub>2</sub> NPs are known to produce reactive oxygen species (ROS) due to its photocatalytic property (Hund-Rinke and Simon, 2006) resulting in lipid peroxidation, damaging DNA, proteins and lipids (Kelly *et al.*, 1998). Increased levels of MDA observed in our study indicate oxidative stress and peroxidation of biological membrane (Fig 5a). Report by Ghosh *et al.* (2010) showed an increase in MDA content in *Allium cepa* as a result of TiO<sub>2</sub> NP treatment. They also

reported geno-toxicity potential of TiO<sub>2</sub> NPs in both plant and human lymphocytes.

Exposure to heavy metals is known to cause water imbalance in the plants, suggesting that proline accumulation induced by metals depends on the development of metal-induced water stress in plant leaves (Chen *et al.*, 2001). In our study, we observed that proline, an osmo-protectant increased with increase in TiO<sub>2</sub> NP concentration, indicating osmotic stress (Fig 5b). As a consequence of the osmotic stress, stomatal limitation was observed (Fig 3) which probably limited the CO<sub>2</sub> fixation and transpiration rate (Fig 3) as reported in this study.

ASA is an important component of the antioxidant defense system in plants serving as a reductant for the peroxidative removal of H<sub>2</sub>O<sub>2</sub> (Noctor and Foyer, 1998). Increase in the non-enzymatic antioxidant, ascorbic acid (ASA) (Fig 6c), observed in this study was probably in order to mitigate, oxidative as well as osmotic stress induced by TiO<sub>2</sub> NPs. Guo *et al.* (2005) reported that ASA decreased the electrolyte leakage in rice roots subjected to Al toxicity and alleviated the inhibition of Al on root growth, indicating that ASA and its precursor increased the resistance of rice seedlings to Al toxicity. They also suggested that ASA decreased the H<sub>2</sub>O<sub>2</sub> and MDA level in rice seedlings exposed to chilling, drought and Al stress. Our results infer that a significant increase in ascorbate content at high TiO<sub>2</sub> NP treatment (Fig 5c) caused a lesser amount of lipid peroxidation (Fig 5a) in *Oryza sativa* var. Jyoti.

## CONCLUSION

Titanium dioxide nanoparticle effect (TiO<sub>2</sub> NPs) on plant growth (root, shoot length and biomass) and photosynthesis was investigated. TiO<sub>2</sub> NPs used in this study showed no effect on seed germination, root and shoot length and light reaction of photosynthesis (Fv/Fm ratio and qP), however, it decreased carbon dioxide assimilation (A), transpiration rate (E) and stomatal conductance (gs) in *Oryza sativa* var. Jyoti at high concentration. TiO<sub>2</sub> NP treatment resulted in an increased MDA, proline and ascorbate content at high concentration indicating oxidative as well as osmotic stress to plants. Overall, TiO<sub>2</sub> NPs caused little or no effect on light reaction of photosynthesis and pigment biosynthesis but led to a phytotoxic effect on the dark reaction of photosynthesis (carbon dioxide fixation).

## Acknowledgements

The authors would like to thank the Department of Science & Technology (DST), New Delhi (SR/SO/PS-63/2009) and UGC- SAP (F. 3-50/2009 (SAP-II) for funding this work. The authors also thank, NIO, Goa for SEM imaging and ICP-OES analysis. The authors are thankful to the Physics Department, Goa University using X-ray Diffractometer and the Centre for Nano Science and Engineering (CeNSE), Indian Institute of Science, Bangalore, India for NP size determination. The authors appreciate the valuable suggestions given by the reviewers.

## References

- Asli, S., Neumann, P.M., 2009. Colloidal suspensions of clay or titanium dioxide nanoparticles can inhibit leaf growth and transpiration via physical effects on root water

- transport. *Plant. Cell Environ.* 32, 577–84. doi:10.1111/j.1365-3040.2009.01952.x
- Bates, L.S., Waldren, R.P., Teare, I.D., 1973. Rapid determination of free proline for water-stress studies. *Plant Soil* 39, 205–207. doi:10.1007/BF00018060
- Baum, A., Hartmann, N.B., Grieger, K., Kusk, K.O., 2008. Ecotoxicity of engineered nanoparticles to aquatic invertebrates: a brief review and recommendations for future toxicity testing. *Ecotoxicology* 17, 387–395. doi:10.1007/s10646-008-0208-y
- Boxall, A. B. A., Chaudhry, Q., Sinclair, C., Jones, A., Aitken, R., Jefferson, B., & Watts, C., 2007. Current and future predicted environmental exposure to engineered nanoparticles. Central Science Laboratory, Department of the Environment and Rural Affairs, London, UK, 89.
- Cai, R., Van, G.M., Aw, P.K., Itoh, K., 2006. Solar-driven self-cleaning coating for a painted surface. *Comptes Rendus Chim.* 9, 829–835. doi:10.1016/j.crci.2005.04.007
- Chen, C.T., Chen, L.-M., Lin, C.C., Kao, C.H., 2001. Regulation of proline accumulation in detached rice leaves exposed to excess copper. *Plant Sci.* 160, 283–290. doi:10.1016/S0168-9452(00)00393-9
- Fang, J., Shan, X., Wen, B., Lin, J., Owens, G., 2009. Stability of titania nanoparticles in soil suspensions and transport in saturated homogeneous soil columns. *Environ. Pollut.* 157, 1101–9. doi:10.1016/j.envpol.2008.11.006
- Farré, M., Gajda-Schranz, K., Kantiani, L., Barceló, D., 2009. Ecotoxicity and analysis of nanomaterials in the aquatic environment. *Anal. Bioanal. Chem.* 393, 81–95. doi:10.1007/s00216-008-2458-1
- Franklin NM, Rogers NJ, Apte SC, Batley GE, Gadd GE, C.P., 2007. Comparative toxicity of nanoparticulate ZnO, bulk ZnO, and ZnCl<sub>2</sub> to a freshwater microalga (*Pseudokirchneriella subcapitata*): the importance of particle solubility. *Env. Sci Technol* 41, 8484–90.
- French, R.A., Jacobson, A.R., Kim, B., Isley, S.L., Penn, R.L., Baveye, P.C., 2009. Influence of Ionic Strength, pH, and Cation Valence on Aggregation Kinetics of Titanium Dioxide Nanoparticles. *Environ. Sci. Technol.* 43, 1354–1359. doi:10.1021/es802628n
- Ghosh, M., Bandyopadhyay, M., Mukherjee, A., 2010. Genotoxicity of titanium dioxide (TiO<sub>2</sub>) nanoparticles at two trophic levels: Plant and human lymphocytes. *Chemosphere* 81, 1253–1262. doi:10.1016/j.chemosphere.2010.09.022
- Guo, Z., Tan, H., Zhu, Z., Lu, S., Zhou, B., 2005. Effect of intermediates on ascorbic acid and oxalate biosynthesis of rice and in relation to its stress resistance. *Plant Physiol. Biochem.* 43, 955–62. doi:10.1016/j.plaphy.2005.08.007
- Guzman, K. a D., Finnegan, M.P., Banfield, J.F., 2006. Influence of surface potential on aggregation and transport of titania nanoparticles. *Environ. Sci. Technol.* 40, 7688–93.
- Hara, T., Sonoda, Y., Iwai, I., 1976. Growth response of cabbage plants to transition elements under water culture conditions. *Soil Sci. Plant Nutr.* 22, 307–315. doi:10.1080/00380768.1976.10432993
- Hong, F., Yang, F., Liu, C., Gao, Q., Wan, Z., Gu, F., Wu, C., Ma, Z., Zhou, J., Yang, P., 2005. Influences of nano-TiO<sub>2</sub> on the chloroplast aging of spinach under light. *Biol. Trace Elem. Res.* 104, 249–60. doi:10.1385/BTER:104:3:249
- Hug, S.J., Sulzberger, B., 1994. In situ Fourier Transform Infrared Spectroscopic Evidence for the Formation of Several Different Surface Complexes of Oxalate on TiO<sub>2</sub> in the Aqueous Phase. *Langmuir* 10, 3587–3597. doi:10.1021/la00022a036
- Hund-Rinke, K., Simon, M., 2006. Ecotoxic Effect of Photocatalytic Active Nanoparticles (TiO<sub>2</sub>) on Algae and Daphnids (8 pp). *Environ. Sci. Pollut. Res. - Int.* 13, 225–232. doi:10.1065/espr2006.06.311
- Kaegi, R., Ulrich, A., Sinnet, B., Vonbank, R., Wichser, A., Zuleeg, S., Simmler, H., Brunner, S., Vonmont, H., Burkhardt, M., Bollner, M., 2008. Synthetic TiO<sub>2</sub> nanoparticle emission from exterior facades into the aquatic environment. *Environ. Pollut.* 156, 233–9. doi:10.1016/j.envpol.2008.08.004
- Kahru, A., Dubourguier, H.-C., Blinova, I., Ivask, A., Kasemets, K., 2008. Biotests and Biosensors for Ecotoxicology of Metal Oxide Nanoparticles: A Minireview. *Sensors* 8, 5153–5170. doi:10.3390/s8085153
- Kampfenkel, K., Vanmontagu, M., Inze, D., 1995. Extraction and Determination of Ascorbate and Dehydroascorbate from Plant Tissue. *Anal. Biochem.* 225, 165–167. doi:10.1006/abio.1995.1127
- Kelly, K.A., Havrilla, C.M., Brady, T.C., Abramo, K.H., Levin, E.D., 1998. Oxidative stress in toxicology: established mammalian and emerging piscine model systems. *Environ. Health Perspect.* 106, 375–84.
- Klaine, S.J., Alvarez, P.J.J., Batley, G.E., Fernandes, T.F., Handy, R.D., Lyon, D.Y., Mahendra, S., McLaughlin, M.J., Lead, J.R., 2008. Nanomaterials in the environment: behavior, fate, bioavailability, and effects. *Environ. Toxicol. Chem.* 27, 1825–1851.
- Kurepa, J., Paunesku, T., Vogt, S., Arora, H., Rabatic, B.M., Lu, J., Wanzer, M.B., Woloschak, G.E., Smalle, J.A., 2010. Uptake and distribution of ultrasmall anatase TiO<sub>2</sub> Alizarin red S nanoconjugates in *Arabidopsis thaliana*. *Nano Lett.* 10, 2296–302. doi:10.1021/nl903518f
- Larue, C., Khodja, H., Herlin-Boime, N., Brisset, F., Flank, a M., Fayard, B., Chaillou, S., Carrière, M., 2011. Investigation of titanium dioxide nanoparticles toxicity and uptake by plants. *J. Phys. Conf. Ser.* 304, 012057. doi:10.1088/1742-6596/304/1/012057
- Lu, CM., Zhang, C.Y.W., Tao, M. X., 2002. Research of the effect of nanometer on germination and growth enhancement of *Glycine max L.* and its mechanism. *Soybean Sci.* 21, 168–172.
- Maynard, A.D., Aitken, R.J., Butz, T., Colvin, V., Donaldson, K., Oberdörster, G., Philbert, M.A., Ryan, J., Seaton, A., Stone, V., Tinkle, S.S., Tran, L., Walker, N.J., Warheit, D.B., 2006. Safe handling of nanotechnology. *Nature* 444, 267–269. doi:10.1038/444267a
- Mu, L., Sprando, R.L., 2010. Application of nanotechnology in cosmetics. *Pharm. Res.* 27, 1746–9. doi:10.1007/s11095-010-0139-1
- Mueller, N.C., Nowack, B., 2008. Exposure modeling of engineered nanoparticles in the environment. *Environ. Sci. Technol.* 42, 4447–53.
- Navarro, E., Baum, A., Behra, R., Hartmann, N.B., Filser, J., Miao, A.-J., Quigg, A., Santschi, P.H., Sigg, L., 2008. Environmental behavior and ecotoxicity of engineered nanoparticles to algae, plants, and fungi. *Ecotoxicology* 17, 372–86. doi:10.1007/s10646-008-0214-0

- Noctor, G., Foyer, C.H., 1998. Ascorbate and glutathione: Keeping Active Oxygen Under Control. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 49, 249–279. doi:10.1146/annurev.arplant.49.1.249
- Ojamäe, L., Aulin, C., Pedersen, H., Käll, P.-O., 2006. IR and quantum-chemical studies of carboxylic acid and glycine adsorption on rutile TiO<sub>2</sub> nanoparticles. *J. Colloid Interface Sci.* 296, 71–78. doi:10.1016/j.jcis.2005.08.037
- Rico, C.M., Majumdar, S., Duarte-Gardea, M., Peralta-Videa, J.R., Gardea-Torresdey, J.L., 2011. Interaction of nanoparticles with edible plants and their possible implications in the food chain. *J. Agric. Food Chem.* 59, 3485–98. doi:10.1021/jf104517j
- Rosenbluh, J., Singh, S.K., Gafni, Y., Graessmann, A., Loyter, A., 2004. Non-endocytic penetration of core histones into petunia protoplasts and cultured cells: a novel mechanism for the introduction of macromolecules into plant cells. *Biochim. Biophys. Acta* 1664, 230–40. doi:10.1016/j.bbamem.2004.06.003
- Sankhalkar, S., Sharma, P.K., 2002. Protection against photooxidative damage provided by enzymatic and non-enzymatic antioxidant system in sorghum seedlings. *Indian J. Exp. Biol.* 40, 1260–1268.
- Schreiber, U., Schliwa, U., Bilger, W., 1986. Continuous recording of photochemical and non-photochemical chlorophyll fluorescence quenching with a new type of modulation fluorometer. *Photosynth. Res.* 10, 51–62. doi:10.1007/BF00024185
- Sharma, P.K., Hall, D.O., 1996. Effect of photoinhibition and temperature on carotenoids in sorghum leaves. *Indian J. Biochem. Biophys.* 33, 471–7.
- Sharma, P.K., Shetye, R., Bhonsle, S., 1997. Effect of supplementary ultraviolet-B radiation on young wheat seedlings. *Curr. Sci.* 72, 400–405.
- Tourinho, P.S., van Gestel, C.A.M., Lofts, S., Svendsen, C., Soares, A.M.V.M., Loureiro, S., 2012. Metal-based nanoparticles in soil: fate, behavior, and effects on soil invertebrates. *Environ. Toxicol. Chem.* 31, 1679–92. doi:10.1002/etc.1880
- Watanabe, T., Misawa, S., Hiradate, S., Osaki, M., 2008. Root mucilage enhances aluminum accumulation in *Melastoma malabathricum*, an aluminum accumulator. *Plant Signal. Behav.* 3, 603–5.
- Wierzbicka, M., Obidzi ska, J., 1998. The effect of lead on seed imbibition and germination in different plant species. *Plant Sci.* 137, 155–171. doi:10.1016/S0168-9452(98)00138-1
- Wiesner, M.R., Lowry, G. V., Alvarez, P., Dionysiou, D., Biswas, P., 2006. Assessing the Risks of Manufactured Nanomaterials. *Environ. Sci. Technol.* 40, 4336–4345. doi:10.1021/es062726m
- Yang, L., Watts, D.J., 2005. Particle surface characteristics may play an important role in phytotoxicity of alumina nanoparticles. *Toxicol. Lett.* 158, 122–32. doi:10.1016/j.toxlet.2005.03.003
- Yuranova, T., Laub, D., Kiwi, J., 2007. Synthesis, activity and characterization of textiles showing self-cleaning activity under daylight irradiation. *Catal. Today* 122, 109–117. doi:10.1016/j.cattod.2007.01.040
- Zheng, L., Hong, F., Lu, S., Liu, C., 2005. Effect of nano-TiO<sub>2</sub> on strength of naturally aged seeds and growth of spinach. *Biol. Trace Elem. Res.* 104, 83–92. doi:10.1385/BTER:104:1:083

\*\*\*\*\*