



International Journal Of
**Recent Scientific
Research**

ISSN: 0976-3031
Volume: 7(5) May -2016

**EFFECT OF SUDARSHAN KRIYA YOGA ON SYMPATHETIC FUNCTION TEST IN
HYPERTENSIVE PATIENT**

Agarwal Nitesh., Bijaraniya Kuldeep., Chaturvedi Mukesh
Kumar and Sharma Sudhir Kumar



THE OFFICIAL PUBLICATION OF
INTERNATIONAL JOURNAL OF RECENT SCIENTIFIC RESEARCH (IJRSR)
<http://www.recentscientific.com/> recentscientific@gmail.com



ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

International Journal of Recent Scientific Research
Vol. 7, Issue, 5, pp. 10909-10912, May, 2016

**International Journal of
Recent Scientific
Research**

Research Article

EFFECT OF SUDARSHAN KRIYA YOGA ON SYMPATHETIC FUNCTION TEST IN HYPERTENSIVE PATIENT

Agarwal Nitesh¹, Bijaraniya Kuldeep², Chaturvedi Mukesh Kumar³
and Sharma Sudhir Kumar⁴

¹Department of Physiology, Government, Medical College, Kota, Rajasthan, India

²Tutor, Department of Physiology, R.N.T. Medical College, Udaipur, Rajasthan, India

³Tutor, Department of Physiology, S.P. Medical College, Bikaner, Rajasthan, India

⁴Medical Officer, Satellite Hospital, Bikaner, Rajasthan, India

ARTICLE INFO

Article History:

Received 15th February, 2016

Received in revised form 21st March, 2016

Accepted 06th April, 2016

Published online 28th May, 2016

Keywords:

Sympathetic functions, Blood pressure, Hand grip test, Sudarshan Kriya.

ABSTRACT

The Sudarshan Kriya is a unique cyclical controlled yogic breathing process developed from ancient Indian science of Yoga. It has potential to integrate body and mind. It is known to eliminate stress, fatigue and negative emotions such as anger, frustration and depression, leaving you calm yet energized, focused yet relaxed. This study evaluated the effects of sudarshan Kriya on autonomic sympathetic variable in hypertensive subjects. In present study two groups each having 35 subjects one is hypertensive and one is normotensive were selected between age 25-75 years of either sex were studied in the morning hours. The patients of hypertension were randomly selected reporting to department of Medicine / Cardiology PBM Hospital, Bikaner. Study group were asked to do Sudarshan Kriya Yoga for three months after proper training under the supervision of expert at The Art of Living Centre Rani Bazar, Bikaner. This study was conducted in the department of physiology Sardar Patel Medical College, Bikaner. Sympathetic non invasive autonomic function test were performed that is hand grip test and postural change test including resting blood pressure. The results were statistically analysed by applying paired "t" test. After the twelve weeks of sudarshan kriya sympathetic function test like orthostatic variation of blood pressure and hand grip test were statistically significantly lower ($p < 0.05$). Reduction was more profound in study group than in control group.

Copyright © Agarwal Nitesh *et al.*, 2016, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Nowadays stress is an absolutely inevitable part of life. The term 'stress' was first employed in the 1930's by the endocrinologist – Hans Selye.¹ Stress is "a physical or psychological stimulus that affects both physical and mental health. Stress causes hormonal changes; biochemical changes various illnesses and psychosomatic diseases. Exposure to chronic stress leads to certain physiological changes in our body resulting in diseased states like hypertension, diabetes, heart attacks etc."²

Hypertension is one of the most important public health problems worldwide. Hypertension is a common disease in industrialized countries, it accounts for 6% of deaths worldwide. It is associated with increased cardiovascular risk which is doubled with 20 and 10 mmHg rise of systolic and diastolic blood pressure respectively. Global epidemic of high blood pressure is expected to shift the burden of disease so that

heart disease will become the most common cause of death worldwide by the year 2025.³ Due to certain physiological disorders the lumen of the arteries constricts, their volume also reduces. This creates increased pressure inside the arteries which is then referred to as "Hypertension". High B.P. may have an hormonal cause among other causes leading to disruption of renin- angiotensin- aldosterone actions, high B.P. may also be as a result of renal disorders. The so called essential hypertension does not show definite correlation to any factor.⁴

Classification of hypertension

The following table is showing the classification of blood pressure levels in adults above age 18 years.⁵

Blood Pressure Classification	Systolic mmHg	Diastolic mmHg
Normal	<120	<80
Pre-hypertension	120-139	80-89
Stage 1 hypertension	140-159	90-99
Stage 2 hypertension	>160	>100
Isolated systolic hypertension	>140	<90

*Corresponding author: Agarwal Nitesh

Department of Physiology, Government, Medical College, Kota, Rajasthan, India

Sudarshan Kriya Yoga

Sudarshan Kriya Yoga (su = proper, darshan = vision, kriya = purifying action) is a unique rhythmical breathing technique, based on ancient vedic tradition.⁶ It is a special package of yoga, meditation and pranayam with “Sudarshan Kriya” introduced by H. H. Sri Sri Ravishankarji.⁷

Sudarshan Kriya

The Sudarshan Kriya is a rhythmical cyclical controlled breathing process with roots in traditional yoga. It has four distinct components.⁸ It consists of slow, medium, and fast cycles of breathing practiced for a total duration of 35 minutes without any relaxation in between. At the end a person is asked to remain in yoga nidra (tranquil state) for about 10 minutes.

The four main SKY breathing techniques are as follows⁹

1. Ujjayi or “Victorious Breath” is sometimes called “Ocean Breath” because the sound created by the gentle contraction of the laryngeal muscles and partial closure of the glottis is reminiscent of the sound of the sea. This slow breath technique (2 to 4 breaths per minute) increases airway resistance during inspiration and expiration and controls airflow so that each phase of the breath cycle can be prolonged to an exact count. The subjective experience is physical and mental calmness with alertness.
2. During Bhastrika or “Bellows Breath” air is rapidly inhaled and forcefully exhaled at a rate of 30 breaths per minute. It engenders excitation followed by calmness.
3. “Om” is chanted three times with very prolonged expiration.
4. Sudarshan Kriya or “Proper Vision by Purifying Action” is an advanced form of cyclical breathing at varying rates—slow, medium, and fast.

Sudarshan Kriya technique oxygenates the cells and rapidly flushes out impurities. Physically, the cells are vitalized; emotionally, one feels a sense of balance and contentment.

MATERIAL AND METHOD

Sample size: In present study two groups each having 35 subjects one is hypertensive and one is control group were selected between age 25-75 years of either sex were studied in the morning hours. The patients of hypertension were randomly selected reporting to department of Medicine / Cardiology PBM Hospital, Bikaner.

Subjects who are hypertensive according to WHO guidelines were examined for the cardiovascular autonomic. Study group and control group were asked to do Sudarshan Kriya Yoga for three months after proper training under the supervision of expert at The Art of Living Centre Rani Bazar, Bikaner. This study was conducted in the department of physiology Sardar Patel Medical College, Bikaner.

Controls: Age-matched normotensive (N=35) were examined for the control values of the cardiovascular autonomic functions.

Exclusion criteria

1. Patients suffering for diabetes mellitus, pulmonary tuberculosis, asthma, chronic bronchitis and no other allergic condition.
2. Absence of any major psychiatric illness. They should not be on any medication of mental illness

Evaluations: The following tests were performed for assessment of sympathetic activity

Resting blood pressure

Blood pressure was recorded with standard sphygmomanometer by auscultatory method. Before recording the blood pressure, subjects were allowed to rest for 5 minutes in a quiet room to reduce the anxiety. The onset of sounds (korkoff’s phase 1) was taken as indicative of systolic blood pressure and disappearance of sound (korkoff’s phase 5) as indicative of diastolic blood pressure.

Lying to standing test (LST)

The subject was asked to rest in a supine position for 5 minutes. The resting BP was recorded. The subject was then asked to stand unaided and remain standing unsupported for 3 minutes. The BP was recorded at 30 seconds and 3 minutes after standing up. The difference between the resting and standing BP levels was calculated.

The fall in systolic BP at 30 seconds on standing noted. A fall of 30 mm Hg or more was defined as abnormal, fall between 11-29 mm Hg as borderline and fall of 10 mm Hg or less was considered normal.

Isometric handgrip test (HGT)

In this test, sustained muscle contraction is measured by a handgrip dynamometer, causes a rise in systolic and diastolic blood pressure and heart rate. The dynamometer is first squeezed to isometric maximum, and then held at 30% maximum for 5 min. if possible, although even 3 minutes may be adequate. Blood pressure was recorded in the non exercising arm thrice at 1-minute interval during the procedure. The maximum reading of the diastolic blood pressure was taken as the final value. Then the rise in diastolic blood pressure was calculated by subtracting resting diastolic blood pressure from this value. A rise in DBP of less than 10 mm Hg was defined as abnormal, 11-15 mm Hg as borderline and 16 mm Hg or more as normal.

Analysis of Observations

Analysis was done by statistical analysis. Students ‘t’ test (two tailed) has been used to find the significance. P=0.05 was considered as statistically significant.

RESULTS

Table 1 Mean age of subjects under study

	Female		Male		Total	
	control	study	control	study	control	Study
Mean	67.51	60.54	63.71	66.07	63.15	66.12
S.D	5.43	7.26	6.65	5.67	7.55	5.63

Table 2 Anthropometric and Sympathetic reactivity test in Control Group in Male

Parameters	Pre- Intervention Post Intervention		p	
	Mean±SD	Mean±SD		
BMI (kg/m ²)	26.21±3.76	26.10±0.143	0.867	
Blood Pressure (mmHg)	Systolic	118.24±7.97	115.72±8.84	0.391
	Diastolic	79.12±7.56	77.92±6.11	0.114
Sympathetic Reactivity Test	LST	6.04±2.65	6.07±0.05	0.0008
	HGT	14.06±2.00	10.31±2.33	0.0007

Table 3 Anthropometric and Sympathetic reactivity test in Control Group in Female

Parameters	Pre- Intervention Post Intervention		p	
	Mean±SD	Mean±SD		
BMI (kg/m ²)	24.21±3.12	22.11±0.113	0.865	
Blood Pressure (mmHg)	Systolic	110.42±11.65	108.72±8.84	0.391
	Diastolic	73.04±5.99	70.92±6.11	0.134
Sympathetic Reactivity Test	LST	6.04±2.45	6.01±0.04	0.0006
	HGT	13.60±2.47	10.02±2.13	0.0007

Table 4 Anthropometric and Sympathetic reactivity test in study group in Male

Parameters	Pre- Intervention Post Intervention		P	
	Mean±SD	Mean±SD		
BMI (kg/m ²)	34.05±5.23	26.13±1.133	0.543	
Blood Pressure (mmHg)	Systolic	145.45±10.14	131.73±10.44	0.643
	Diastolic	94.04±6.71	82.92±5.17	0.342
Sympathetic Reactivity Test	LST	7.32±2.15	6.01±0.06	0.0008
	HGT	12.60±3.56	9.04±2.23	0.0007

Table 5 Anthropometric and Sympathetic Reactivity test in study group (Hypertensive) in Female

Parameters	Pre- Intervention Post Intervention		p	
	Mean±SD	Mean±SD		
BMI (kg/m ²)	31.65±5.39	25.11±1.23	0.623	
Blood Pressure (mmHg)	Systolic	138.47±10.14	130.75±10.11	0.50
	Diastolic	91.04±6.51	82.67±5.31	0.42
Sympathetic Reactivity Test	LST	7.32±2.15	6.01±0.06	0.0007
	HGT	11.60±3.46	9.56±2.21	0.0007

The mean age of participants was 63.15±7.55 in control group and in study group it was 66.12±5.63 years (mean ± standard deviation) (Table-1)

Data presented in above table 2,3,4,5 shows that orthostatic variation of blood pressure and hand grip test were statistically significantly lower (p<0.05), in both the male and female group of study group as compared to the male and female group of controls. In both the group resting blood pressure were lower but not statistically significantly (p>0.5).

DISCUSSION

Sudarshan Kriya incorporates specific natural rhythms of the breath which harmonize the body, mind and emotions. This unique breathing technique eliminates stress, fatigue and negative emotions such as anger, frustration and depression, leaving you calm yet energized, focused yet relaxed.⁹

Tests assessing autonomic function are based on evaluation of the cardiovascular reflexes triggered by performing specific provocative manoeuvres. Stimuli that raise blood pressure responses to orthostatic testing are in a large part a reflection of sympathetic activity.

Our findings are consistent to Somwanshi *et al* in 2013 who found highly significant improvement in cardio respiratory parameters in healthy subjects after 3 months of Sudarshan Kriya Yoga practices.¹⁰

Effect of SKY in Reducing blood pressure and heart rate

Sustained stimulation of sympathetic nervous (fight and flight response) system increases blood pressure. It is hypothesized that the different cyclical rhythms of Sudarshan Kriya create a variety of vagal, thalamic and cortical effects. During SKY, a sequence of breathing techniques of different frequencies, intensities, lengths, and with end-inspiratory and end-expiratory holds creates varied stimuli from multiple visceral afferents, sensory receptors, and baroreceptors.¹¹

These influence stimulates vagus nerves which in turn induces physiologic changes in organs, glands, and ascending fibers to thalamic generators, the limbic system and cortical areas.¹²

Resistive loading created during ujjayi breathing send afferent input via vagal & spinal sources arising from lung & chest wall structures to parabrachial nucleus (PBN) & locus coeruleus¹³ which also receives projections from NTS feeding back through dorsomedial nucleus to vagal efferents which in turn slows heart rate by increasing parasympathetic & decreasing sympathetic input to SA node¹⁰ thus decreasing the heart rate. During prolonged voluntary expiration intra-thoracic pressure increases and blood from the lungs is squeezed into the heart leading to an increase in stroke volume; baro-receptors in carotid sinus experiences more pressure and discharge more. The increased baroreceptor discharge inhibit the tonic discharge of the vasoconstrictor nerves and excites the vagus innervations of the heart producing vasodilatation a drop in systolic blood pressure and bradycardia.¹⁴

Bradycardia itself results in fall in systolic blood pressure. Diastolic blood pressure depends upon peripheral vascular resistance & lung inflation has been known to decrease systemic vascular resistance.¹⁵ this response is initiated by pulmonary stretch receptors, which bring about withdrawal of sympathetic tone in the skeletal muscle blood vessels, leading to widespread vasodilatations, thus causing a decrease in peripheral resistance¹⁶ and decreasing the diastolic blood pressure. Yoga on long duration affects hypothalamus and brings about decrease in systolic blood pressure through its influence on vasomotor center which leads to decrease in sympathetic tone and peripheral resistance, thus reducing diastolic blood pressure.¹⁷

SKY practice of ‘ujjayi’ shifts parasympathetic dominance via vagal stimulation from somatosensory afferents in the glottis, pharynx, lungs and abdominal viscera. Thus SKY strengthen, balance, stabilize autonomic nervous system functions and establishes parasympathetic dominance with decreased sympathetic drive.¹¹

SKY showed statistically significant decrease after 12 weeks of practice in the values of heart rate, systolic blood pressure and diastolic blood pressure in healthy subjects. The mechanism involved is by creating balance in autonomic nervous system functions by parasympathetic dominance and decreased sympathetic drive.^{18,10}

By voluntarily controlling breathing patterns, it is possible to influence autonomic nervous system functions, including heart rate variability and cardiac vagal tone.^{19,20}

Thus by establishing parasympathetic dominance SKY is beneficial to decrease blood pressure and heart rate in patients of hypertension.

Our study showed that the systolic and diastolic blood pressure response to hand grip test was significantly lower in both the male and female group of control and study groups. ($p < 0.05$)

Our findings are in consistent with Akhter et al,²¹ who conducted a cross sectional study which showed that mean values of resting heart rate, resting systolic and diastolic blood pressure were significantly ($p < 0.05$) higher and both the sympathetic nerve function parameters (blood pressure response to hand grip and blood pressure response to standing) were significantly ($p > 0.05$) lower in hypertensive compared to those of normotensive control subjects. So it can be revealed that sympathetic nerve function may be reduced in hypertension.

Piccirillo et al²², Peterson et al²³, also stated the same that sympathetic nerve function are reduced in hypertension.

CONCLUSION

In our study the sympathetic function tests showed significant ($p < 0.05$) decrease in both the hypertensive and normotensive individual. So we can say Sudarshan Kriya Yoga can be used as an adjunct with diet and medicines in management of hypertension and anxiety.

SKY practices are cost-effective, well-tolerated tools that can be easily integrated into diverse community care models. SKY relieves stress and develops an individual's mind–body–spirit so that they can be happier, healthier, and possibly even longer lived.

In the competitive modern world, in which stress and anxiety are part of everyday life, adding a time-honored, evidence-based breathing program like SKY may facilitate a healthy life. So everyone should incorporate SKY into their life.

Reference

- Hans Selye. The stresses of life, New York, MC Graw Hill.1956; 523-67.
- Christensen NJ, Jensen FW. Effect of psychological stress and age on plasma norepinephrine levels. A review Psychosomatic Medicine. 1994; 56: 77:83.
- Ezzati M, Lopez AD, Rodegrs A. Selected major risk factors and global and regional burden of disease. Lancet 2002; 360:1347-60.
- Http://www.Yoga point. Hypertension & yoga – by. Sujit Chandradratreyan.
- Kotchen TA. Hypertensive vascular disease. In: Harrison's Principles of Internal Medicine. Fauci, Braunwald, Kasper, Hauser, Longo, Jameson, Loscalzo(eds). 17th edition. 2008; II: pp1549-62.
- Vedarmurthachar. The role of Sudarshan Kriya on mental health. International symposium on YoGism.2010; 32-34.
- Vedamurthachar A, Bijoor A R, Agte V, Reddy S, Lakshmi B. Short term effect of Sudarshan Kriya yoga on lipid and hormone profile of type 2 diabetic patients. *Research Journal of Chemical Science*. 2011; 1(9): 83-86.
- Zope SA, Zope RA. Sudarshan Kriya yoga: breathing for health. *Int J Yoga*. 2013; 6: 4-10.
- Brown RP, Gerbarg PL. Sudarshan Kriya yogic breathing in the treatment of stress, anxiety, and depression: Part II-clinical applications and guidelines. *J Altern Complement Med*. 2005; 11: 711–17.
- Somwanshi S, Handergulleb SM, Adgonakar BD, Kolpe D.V. Effect of Sudarshan Kriya Yoga on cardiorespiratory parameters. *Int J Recent Trends in Science and Technology*.2013; 8(1):62-66.
- Brown RP, Gerbarg PL. Sudarshan Kriya yogic breathing in the treatment of stress, anxiety, and depression: Part I-neurophysiologic model. *J Altern Complement Med*. 2005; 11: 189–201.
- Porges SW. The vagus a mediator of behavioral and visceral features associated with autism. In: Bauman ML, Kemper TL, eds. *The Neurobiology of autism*. Baltimore: Johns Hopkins University Press, 2004.
- Gozal D, Omidvar O, Kirlaw KA, Hathout GM, Hamilton R et al. Identification of human brain regions underlying responses to resistive inspratory loading with functional magnetic resonance imageing. *Proc Natl Acad Sci USA*. 1995; 92: 6607-11.
- Ganong WF. Cardiovascular regulatory mechanism. In: *Review of Medical Physiology*. 22nd Edition. 2005; pp 597-610.
- Hainsworh R. Circulatory responses from lung inflation in anaesthetized dogs. *Amer J Physiol*. 1974; 226: 247-55.
- Daly M, De B, Robinson BH. An analysis of the reflex systemic vasodilator response elicited by lung inflation in dog. *J Physiol London*. 1968; 195:387-406.
- Bhaskar R.J. Effect of yoga on cardiovascular system in subjects above 40 years. *Indian J Physiol Pharmacol*. 2003; 47(2)202-206.
- Deepak K K. The role of autonomic nervous system in rapid breathing practices. *Proceedings: Science of breath. International Symposium on Sudarshan Kriya, Pranayam and Consciousness, New Delhi, All India Institute of Medical Sciences*.2002;43-46.
- Fokkema DS. The psychobiology of strained breathing and its cardiovascular implications: A functional system review. *Psychophysiology*. 1999; 36: 164–75.
- Sovik R. The science of breathing-The yogic view. *Progr Brain Res*. 2000; 122: 491-505.
- Akhter S, Begum N, Ferdousi S, Begum S, Ali T. Sympathetic Nerve Function status in obesity. *J Bangladesh Soc Physiol*. 2010;5(1):34-39.
- Piccirillo G, Vetta F, Fimognari FL, et al. Power spectral analysis of heart rate variability in obese subjects: evidence of decreased cardiac sympathetic responsiveness. *Int J Obes Relat Metab Disord J Int Assoc Study Obes*. 1996;20(9):825-829.
- Peterson HR, Rothschild M, Weinberg CR, Fell RD, McLeish KR, Pfeifer MA. Body fat and the activity of the autonomic nervous system. *N Engl J Med*. 1988;318(17):1077-1083.

T.SSN 0976-3031



9 770976 303009 >