



ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

International Journal of Recent Scientific Research
Vol. 7, Issue, 12, pp. 14873-14876, December, 2016

**International Journal of
Recent Scientific
Research**

Research Article

EFFECT OF NITROGEN, PHOSPHORUS AND POTASSIUM LEVELS ON LINSEED (*LINUM USITATISSIMUM L.*)

Patil Vidya U

Department of Chemistry, Gokhale Education Society's HPT Arts and RYK Science College,
Nasik -422005, Maharashtra, India

ARTICLE INFO

Article History:

Received 10th September, 2016
Received in revised form 14th
October, 2016
Accepted 08th November, 2016
Published online 28th December, 2016

Key Words:

Linseed, Nitrogen, Phosphorus, Potassium.

ABSTRACT

The present investigation was conducted during *rabi* 2015- 2016 at Chincholi near Nashik. The experiment was laid out in randomized block design (Factorial) with three level each of nitrogen ($N_1: 40, N_2: 60, N_3: 80 \text{ kg ha}^{-1}$), phosphorus ($P_1: 15, P_2: 30, P_3: 45 \text{ kg ha}^{-1}$) and potassium ($K_1: 0, K_2: 15, K_3: 30 \text{ kg ha}^{-1}$) replicated twice. Chemical studies the nutrient concentrations as influenced by different treatments viz., N, P and K content (%) in seed and straw, N, P and K uptake by seed, straw and crop (kg ha^{-1}) and available N, P and K in soil (kg ha^{-1}) after harvest of crop.

The mean concentration of nitrogen, phosphorus, potassium in linseed seed were 4.63, 0.84 and 0.85 per cent, while in straw 1.06, 0.54 and 1.03 per cent, respectively. Total N, P and K uptake (kg ha^{-1}) by seed, straw and crop were increased significantly due to application of 80:45:30 NPK kg ha^{-1} and Available N, P and K (kg ha^{-1}) in soil after harvest was higher at application of 80 kg ha^{-1} nitrogen, 45 kg ha^{-1} phosphorus and 30 kg ha^{-1} potassium over lower levels of fertilizer application.

Copyright © Patil Vidya U., 2016, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Linseed is one of the important *rabi* oilseed crop of India. Linseed (*Linum usitatissimum L.*) is one of the oldest oilseed crop cultivated for its seed and fiber. It is an industrial crop cultivated for its seeds, fibers and oil. All parts of the plant have extensive and varied uses.

The oil content of seed varies from 32 to 46 per cent. Linseed oil is found to be comprised of 5 major fatty acids, viz., palmitic, stearic, oleic, linoleic and linolenic.

The linseed oil primarily goes to industries for manufacture of paints, varnish, oil cloth, linoleum and pad ink. Linseed oil possesses a very healthy fatty acid profile, particularly, Omega-3 (Alpha Linolenic Acid), the richest source only in linseed (58 per cent). Every part of the linseed plant is utilized commercially either directly or after processing ALA provides beneficial effects in numerous clinical conditions such as, cardiovascular disease, inflammatory disorders, immune function and cancer etc.

Linseed responds to N, P and K nutrients very well. The beneficial effect of N, P and K in increasing growth attributes of linseed was reported by several workers (Jain *et al.*, 1989). Studies on direct effect of N, P and K fertilization on the uptake of nutrients by crops have been found to be helpful in economizing fertilizer use.

MATERIALS AND METHODS

An experiment was conducted at *rabi* 2015- 2016 at Chincholi near Nashik. The soil of experiment field was clay loam in texture, normal in reaction (pH 7.6) with low available nitrogen (178.4 kg ha^{-1}), medium available phosphorus (30.46 kg ha^{-1}) and high available potassium ($462.10 \text{ kg ha}^{-1}$).

The experiment was laid out in randomized block design (Factorial) with three level each of nitrogen ($N_1: 40, N_2: 60, N_3: 80 \text{ kg ha}^{-1}$), phosphorus ($P_1: 15, P_2: 30, P_3: 45 \text{ kg ha}^{-1}$) and potassium ($K_1: 0, K_2: 15, K_3: 30 \text{ kg ha}^{-1}$) replicated twice. The gross and net plot sizes were $4.00 \times 3.00 \text{ m}^2$ and $3.50 \times 2.40 \text{ m}^2$, respectively. The linseed variety PKVNL- 260 was sown on 5th December, 2015 with spacing of $30 \times 5 \text{ cm}^2$.

RESULTS AND DISCUSSION

Chemical studies

Nutrient concentration

The nutrient concentrations as influenced by different treatments. The mean concentration of nitrogen, phosphorus, potassium in linseed seed were 4.63, 0.84 and 0.85 per cent, while in straw 1.06, 0.54 and 1.03 per cent, respectively.

Effect of nitrogen levels

The effect of N levels viz., 40, 60 and 80 kg ha^{-1} on NPK content in seed and straw were statistically non significant.

*Corresponding author: Patil Vidya U

Department of Chemistry, Gokhale Education Society's HPT Arts and RYK Science College, Nasik-422005, Maharashtra, India

Numerically, the nutrient concentration in seed and straw were not showed considerable differences.

Table 1 Total N, P and K content (%) in seed and straw as influenced by different treatments

Treatment	NPK content in seed (%)			NPK content in straw (%)		
	N	P	K	N	P	K
Nitrogen levels (kg ha⁻¹)						
N ₁ : 40	4.85	0.99	1.01	1.37	0.70	1.21
N ₂ : 60	5.04	1.04	1.06	1.49	0.75	1.24
N ₃ : 80	5.21	1.07	1.09	1.54	0.77	1.26
S.E. m ±	0.008	0.003	0.001	0.001	0.001	0.001
C.D. at 5%	NS	NS	NS	NS	NS	NS
Phosphorus levels (kg ha⁻¹)						
P ₁ : 15	4.99	1.02	1.04	1.24	0.73	1.22
P ₂ : 30	5.03	1.04	1.05	1.26	0.74	1.23
P ₃ : 45	5.07	1.05	1.06	1.29	0.75	1.25
S.E. m ±	0.008	0.003	0.001	0.001	0.001	0.001
C.D. at 5%	NS	NS	NS	NS	NS	NS
Potassium levels (kg ha⁻¹)						
K ₁ : 0	5.02	1.03	1.05	1.25	0.73	1.23
K ₂ : 15	5.03	1.04	1.05	1.26	0.74	1.24
K ₃ : 30	5.04	1.04	1.05	1.27	0.74	1.24
S.E. m ±	0.008	0.003	0.001	0.001	0.001	0.001
C.D. at 5%	NS	NS	NS	NS	NS	NS
Interaction						
N x P	NS	NS	NS	NS	NS	NS
N x K	NS	NS	NS	NS	NS	NS
P x K	NS	NS	NS	NS	NS	NS
N x P x K	NS	NS	NS	NS	NS	NS
General Mean	5.03	1.04	1.05	1.26	0.74	1.23

Effect of phosphorus levels

The P levels viz., 15, 30 and 45 kg ha⁻¹ were statistically non significant in seed and straw. Numerically the nutrient concentration was not showed considerable difference in nutrient concentration of seed and straw.

Effect of potassium levels

The effect of K levels viz., 0, 15 and 30 kg ha⁻¹ were statistically non significant in seed and straw. Numerically, the nutrient concentration of seed and straw was not showed considerable differences.

Table 2 Total N, P and K uptake (kg ha⁻¹) by seed, straw and crop as influenced by different treatments

Treatment	NPK uptake by seed			NPK uptake by straw			Total NPK uptake by crop		
	N	P	K	N	P	K	N	P	K
Nitrogen levels (kg ha⁻¹)									
N ₁ : 40	49.48	8.91	9.13	23.43	12.07	24.28	72.91	20.98	33.13
N ₂ : 60	71.06	12.99	13.29	30.54	15.55	29.22	102.20	28.54	42.51
N ₃ : 80	83.27	15.19	15.53	35.30	17.65	32.65	118.17	32.84	48.18
S.E. m ±	0.93	0.26	0.24	0.48	0.16	0.41	0.94	0.28	0.60
C.D. at 5%	2.80	0.78	0.73	1.44	0.48	1.25	2.84	0.86	1.80
Phosphorus levels (kg ha⁻¹)									
P ₁ : 15	58.64	10.59	10.85	26.44	13.47	25.74	85.08	24.06	36.59
P ₂ : 30	67.21	12.32	12.46	29.68	11.12	28.65	96.89	27.44	41.11
P ₃ : 45	76.89	14.12	14.28	32.66	16.48	31.28	109.55	30.60	45.56
S.E. m ±	0.93	0.26	0.24	0.48	0.16	0.41	0.94	0.28	0.60
C.D. at 5%	2.80	0.78	0.73	1.44	0.48	1.25	2.84	0.86	1.80
Potassium levels (kg ha⁻¹)									
K ₁ : 0	57.22	10.40	10.24	26.65	13.45	25.95	83.87	23.65	36.19
K ₂ : 15	66.84	12.25	13.19	30.08	15.32	29.32	96.92	27.57	42.51
K ₃ : 30	77.99	14.27	14.44	31.64	15.96	30.56	109.63	30.23	45.00
S.E. m ±	0.93	0.26	0.24	0.48	0.16	0.41	0.94	0.28	0.60
C.D. at 5%	2.80	0.78	0.73	1.44	0.48	1.25	2.84	0.86	1.80
Interaction									
N x P	NS	NS	NS	NS	NS	NS	NS	NS	NS
N x K	NS	NS	NS	NS	NS	NS	NS	NS	NS
P x K	NS	NS	NS	NS	NS	NS	NS	NS	NS
N x P x K	NS	NS	NS	NS	NS	NS	NS	NS	NS
General Mean	67.62	12.44	12.60	29.60	15.00	28.61	97.22	27.34	41.19

Interaction effects

The interactions effect nitrogen, phosphorus and potassium levels were statistically non significant for nutrient concentration in seed and straw by linseed.

Total nutrient uptake

The data on mean total uptake of nitrogen, phosphorus and potassium by linseed plant at harvest as influenced by the different treatments.

The mean total uptake of nitrogen, phosphorus and potassium were 97.22, 27.34 and 41.19 kg ha⁻¹, respectively.

Effect of nitrogen levels

The total uptake of nitrogen, phosphorus and potassium was influenced significantly by N levels. Significantly higher total nitrogen, phosphorus and potassium uptake i.e.83.27,15.19, 15.53 and 35.30, 17.65, 32.65 in seed and straw respectively were registered in N level of 80 kg ha⁻¹ over rest of N levels. The higher nutrients uptake with N level 80 kg ha⁻¹ was might be due to higher N availability.

The similar results were reported by Chourasia *et al.* (1992), Dixit and Sharma (1993), Vashishtha, (1993), Agrawal *et al.* (1994), Jain *et al.* (1997) and Sharma and Hunsingi, (1996).

Effect of phosphorus levels

The total uptake of nitrogen, phosphorus and potassium was influenced significantly by P levels. Significantly the highest total nitrogen, phosphorus and potassium uptake were registered with 45 kg ha⁻¹ i.e.76.89,14.12, 14.28 and 32.66, 16.48, 31.28 in seed and straw respectively .Whereas, it was significantly minimum with 15 kg ha⁻¹ i.e.58.64,10.59,10.85 and 26.44,13.47, 25.74 in seed and straw respectively. The higher nutrients uptake in P level 45 kg ha⁻¹ was might be due to higher P availability.

The similar results were reported by Vashishtha, (1993) and Jain *et al.* (1997).

Effect of potassium levels

The total uptake of nitrogen, phosphorus and potassium was influenced significantly by K levels. Total nitrogen, phosphorus and potassium uptake by plant were increased progressively with each successive increase in potash levels. Increased in nutrients uptake by plants due to higher levels of potash might be due to higher K availability. Similar results were observed by Dixit and Sharma (1993), Agrawal *et al.* 1994 and Jain *et al.* (1997).

Interaction effects

The interactions effect of nitrogen, phosphorus and potassium levels were statistically non significant for nutrient removal by seed, straw and total uptake by linseed.

Nutrient status of soil

The data in respect of available of nitrogen, phosphorus and potassium in soil after harvest as influenced by the different treatments are presented in Table 3. The mean total available nitrogen, phosphorus and potassium after harvest were 132.34, 24.21 and 428.70 kg ha⁻¹, respectively.

Table 3 Available N, P and K (kg ha⁻¹) in soil after harvest as influenced by different treatments

Treatment	Nitrogen	Phosphorus	Potassium
Nitrogen levels (kg ha⁻¹)			
N ₁ : 40	129.04	19.82	422.51
N ₂ : 60	133.07	23.02	427.81
N ₃ : 80	138.73	26.68	437.83
S.E. m ±	0.42	0.31	0.53
C.D. at 5%	1.27	0.94	1.60
Phosphorus levels (kg ha⁻¹)			
P ₁ : 15	127.56	21.12	417.93
P ₂ : 30	134.75	24.67	429.01
P ₃ : 45	141.09	27.68	439.79
S.E. m ±	0.42	0.31	0.53
C.D. at 5%	1.27	0.94	1.60
Potassium levels (kg ha⁻¹)			
K ₁ : 0	124.04	22.42	417.12
K ₂ : 15	128.07	23.73	427.42
K ₃ : 30	134.73	28.82	438.88
S.E. m ±	0.42	0.31	0.53
C.D. at 5%	1.27	0.94	1.60
Interaction			
N x P	NS	NS	NS
N x K	NS	NS	NS
P x K	NS	NS	NS
N x P x K	NS	NS	NS
General Mean	132.34	24.21	428.70

Effect of nitrogen levels

The nutrient status of soil was influenced significantly due to N levels. The available residual nitrogen, phosphorus and potassium were registered significantly maximum in N level 80 kg ha⁻¹. Whereas, these were significantly lower with 40 kg N ha⁻¹. This might be due to more availability of nitrogen, phosphorus and potassium in soil at higher level of nitrogen i.e. 80 kg ha⁻¹ and reflects in more uptake and higher yield.

Effect of phosphorus levels

The nutrient status of soil was influenced significantly due to P levels. The available residual nitrogen, phosphorus and potassium were registered significantly maximum in P level of 45 kg ha⁻¹. Whereas, significantly lowest available nitrogen, phosphorus and potassium in soil were recorded in P level 15 kg ha⁻¹. might be due to more availability of nitrogen,

phosphorus and potassium in soil at P level of 45 kg ha⁻¹ which reflects in more uptake and higher yield.

Effect of potassium levels

The nutrient status of soil was influenced significantly due to K levels. The available residual nitrogen, phosphorus and potassium were registered significantly maximum in K level 30 kg ha⁻¹. Whereas, significantly lower nitrogen, phosphorus and potassium content in soil were recorded in K levels of 0 kg ha⁻¹. This difference might be due to more availability of nitrogen, phosphorus and potassium under higher level of potash resulted in more uptake and higher yield.

Interaction effects

The interaction effect among nitrogen, phosphorus and potassium levels were statistically non significant for available residual soil nitrogen, phosphorus and potassium.

CONCLUSION

The N level 80 kg ha⁻¹ recorded significant the maximum uptake of total nitrogen (118.17 kg ha⁻¹), phosphorus (32.84 kg ha⁻¹) and potassium (48.18 kg ha⁻¹) by linseed.

After harvest of crop available soil nitrogen (138.73 kg ha⁻¹), phosphorus (26.68 kg ha⁻¹) and potassium (437.83 kg ha⁻¹) were significantly more with 80 kg ha⁻¹ N level as compared with 60 kg ha⁻¹ and 40 kg ha⁻¹.

The P level 45 kg ha⁻¹ recorded the significantly maximum uptake of total nitrogen (109.55 kg ha⁻¹), phosphorus (30.60 kg ha⁻¹) and potassium (45.56 kg ha⁻¹) by linseed.

After harvest of crop available soil nitrogen (141.09 kg ha⁻¹), phosphorus (27.68 kg ha⁻¹) and potassium (439.79 kg ha⁻¹) were significantly higher with 45 kg ha⁻¹ over with 30 and 15 kg ha⁻¹.

The K level of 30 kg ha⁻¹ recorded the significantly maximum uptake of total nitrogen (109.63 kg ha⁻¹), phosphorus (30.23 kg ha⁻¹) and potassium (45.00 kg ha⁻¹) by linseed.

After harvest of crop available soil nitrogen (134.73 kg ha⁻¹), phosphorus (28.82 kg ha⁻¹) and potassium (438.88 kg ha⁻¹) were significantly more with 30 kg ha⁻¹ K level over 15 and 0 kg ha⁻¹.

References

- Agrawal, K.K., Tiwari, J.P. and Jain, K.K. 1994. Influence of irrigation and fertility levels on utilization and oil content in linseed. (*Linum usitatissimum* L.). *J. Oilseeds Res.* 11 (1): 81-83.
- Chourasia, S.K., Chourasia, S.C. and Namdeo, K.N. 1992. Nitrogen and sulphur uptake by different parts of linseed plant (*Linum usitatissimum* L.) fertilized with these nutrients. *Crop Res. Hisar.* 5: Supp., 65-73.
- Dixit, S.P. and Sharma, P.K. 1993. Effect of lime and potassium on yield and uptake of nutrients in wheat - soybean - linseed cropping sequence in an acid alfisol. *Indian J. Agric. Sci.* 63 (6): 333-339.
- Jain, V.K., Chauhan, Y.S., Khandekar, Sharma, R.P., and Yadav, M.S. 1989. Effect of nitrogen and phosphorus on growth and yield of linseed (*Linum usitatissimum* L.). *Indian J. Agron.* 34 (1): 122-124.

- Jain, N.K., Agrawal, K.K. and Jain, K.K. 1997. Effect of irrigation, fertility and weed control on the availability of nutrients to linseed (*Linum usitatissimum L.*) and removal of nutrients by weeds. *J. Oilseeds Res.* 14 (1): 59-61.
- Sharma, A. and Hunsingi, G. 1996. Influence of genotypes, spacing and nitrogen application on yield and nutrient uptake of linseed. *J. Maharashtra Agric. Univ.*, 21 (1): 137.
- Vashishtha, R.P. 1993. Influence of nitrogen and phosphorus application on the yield and oil content of linseed (*Linum usitatissimum L.*). *Indian J. Agron.* 38 (1): 64-67.

How to cite this article:

Patil Vidya U.2016, Effect of Nitrogen, Phosphorus And Potassium Levels on Linseed (*Linum Usitatissimum L.*). *Int J Recent Sci Res.* 7(12), pp. 14873-14876.