



ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

International Journal of Recent Scientific Research  
Vol. 8, Issue, 3, pp. 15909-15912, March, 2017

**International Journal of  
Recent Scientific  
Research**

## Research Article

### PREVALENCE OF VITAMIN D DEFICIENCY AMONG MENOPAUSAL WOMEN AND ITS CORRELATION WITH DIABETES: A CROSS SECTIONAL DATA

Shylla Mir and Samiya Mufti

Department of Obstetrics and Gynaecology Government Medical College, Srinagar, J&K, India

DOI: <http://dx.doi.org/10.24327/ijrsr.2017.0803.0031>

#### ARTICLE INFO

##### Article History:

Received 06<sup>th</sup> December, 2015  
Received in revised form 14<sup>th</sup>  
January, 2017  
Accepted 23<sup>rd</sup> February, 2017  
Published online 28<sup>th</sup> March, 2017

##### Key Words:

Chronic diseases, deficiency, diabetes,  
postmenopausal women, vitamin D.

#### ABSTRACT

**Aim and objective:** To evaluate prevalence of Vitamin D deficiency and establish any correlation between diabetes and vitamin D deficiency among postmenopausal women.

**Materials and Methods:** The 25 – hydroxy vitamin D [25 (OH)D] concentrations were measured by competitive in- vitro quantitative immunoassay. The subjects were classified as vitamin – D-deficient, insufficient or sufficient on the bases of 25 (OH) D concentrations of <20ng /ml, 20-30ng /ml or > 30ng /ml respectively. The apparently normal postmenopausal women (PMW) were subjected to fasting between vitamin D deficiency and diabetes.

**Results:** Vitamin D deficiency was observed in 53.35% of the population, 19.48% had insufficiency and 26.83% had adequate Vitamin D levels. In 12.14% of the study population, fasting blood sugar levels were between the normal range which is 70-110 mg/ dl. Correlation between raised blood sugar levels and Vitamin D deficiency among PMW was non-significant (P =0.324).

**Conclusion:-** High prevalence of Vitamin D deficiency exists among apparently healthy. Indian PMW. However the current study failed to show any statistical correlation between Vitamin D deficiency and existence of diabetes, which may be due to small sample size.

Copyright © Shylla Mir and Samiya Mufti, 2017, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

#### INTRODUCTION

Vitamin D deficiency is a worldwide health problem, with a range of prevalence of 70% - 100% in the general population.<sup>1</sup> Among Indian population, a prevalence of 50-90% has been reported.<sup>2</sup> Most of the researchers have uniformly reported a high prevalence of vitamin D deficiency, in various groups of the population such as school children, adolescents and pregnant women.<sup>3,4,5.</sup>

There are studies which report high prevalence of Vitamin D deficiency among postmenopausal women (PMW) from western world and India but data still remain scanty<sup>6,7,8,9,10.</sup>

Secondly, Vitamin D deficiency and diabetes mellitus are widely prevalent diseases during menopause. Furthermore, few epidemiological studies have shown an association of vitamin D deficiency and increased risk of chronic diseases like cancer, cardiovascular disease, type 1 and 2 diabetes (T1D, T2D) suggesting that vitamin D deficiency not only affects musculoskeletal health but also affects a wide range of acute and chronic diseases<sup>11.</sup>

#### MATERIALS AND METHODS

This cross-sectional, one year analysis, from June 2015- June 2016 was carried out among apparently healthy postmenopausal subjects (with one year of menstruation cessation) without known history of diabetes, thyroid, renal or hepatic disease or malignancy. Patients with history of surgery, hospitalization, or major medical illness within the past 1 year were excluded from the study patients on hormone replacement therapy, glucocorticoids, bisphosphonates, teriparatide and other drugs affecting bone metabolism were excluded as well. Surgical and other iatrogenic causes of menopause were also excluded from the current study.

Intake of conventional calcium / vitamin D supplements was not considered an exclusion criterion. All subjects were enrolled after taking a written informed consent. The 25 – hydroxy vitamin D [25 (OH) D] concentrations were measured by competitive in-vitro quantitative immunoassay in human serum and Cobas e analyses (Roche) Kit. Inter – assay Coefficient of Variability (CV) was 9.9% and intra-assay CV was 5.7% in the current analysis.

The minimal detectable limit of the 25 (OH) D assays was 1.5ng/ ml. The subjects were classified as vitamin D- deficient

\*Corresponding author: *Shylla Mir*

Department of Obstetrics and Gynaecology Government Medical College, Srinagar, J&K, India

or sufficient on the basis of 25 (OH) D concentrations of < 20ng/ml, to recent consensus.<sup>12,13</sup>

Subjects were included in a cross - sectional manner from general population. The apparently normal PMW were subjected to fasting blood glucose level tests and vitamin D levels to analyse any correlation between the two states. After inclusion in the study they were called on next day for fasting blood samples. Blood glucose levels were estimated by glucose oxidize method and normal reference range was defined as 70 – 110 mg/dl.

Fasting blood was collected and 2ml serum prepared and collected using standard sampling tubes containing separating gel before subjecting them for analysis for vitamin D levels. The data was analysed by one laboratory after standardization.

**Statistical Analysis**

The data was categorized as mean±SD and N (%) and correlation between vitamin D and diabetes was established by using Pearson coefficient correlation test. The P value < 0.05 was considered significant.

**RESULTS**

Total of 312 PMW participated in the current study. Mean age of study population was 56.22 with mean±SD. Number of menopausal symptoms were 3.170±1.16. The current study is depicted in table 1.

Mean vitamin D levels of the study population was 26.86ng/ml and mean fasting blood glucose levels was 134.52±17.56 mg%. While evaluating the vitamin D status of the study population, 53.35% of the population had vitamin D deficiency, 19.48% had in sufficiency and and 26.83% had adequate vitamin D levels (Table 1)

Out of the study population, 12.14% had fasting blood glucose > 110 mg/dl and rest of the population was between the normal range (70-110 mg/dl). The age wise distribution of the population and their vitamin D levels and fasting blood glucose levels are depicted in the correlation between raised blood glucose (Table 2) levels and vitamin D deficiency among PMW was non-significant [(P = 0.324). Table 3].

**Table 1: Profile of study population**

- Study parameters  
 N = 312  
 Mean Age of study population = 56.22  
 Urban Rural = 213 (68.05%): 100 (31.94%)  
 Mean number of symptoms = 3. 170 ± 1.16  
 Most common symptoms
- Fatigue, lack of energy, rheumatic pain = 38%
  - Cold hand and feet = 18%
  - Urogenital symptoms = 18%
  - Cold sweats, weight gain, irritability and nervousness =13%
  - Palpitation of heart, excitable/anxiety/Insomnia = 12%
  - No symptoms = 32%
- Mean vitamin D levels of total population = 26.86ng/ml  
 Mean fasting blood glucose levels ± SD of total population – 134.52 ± 17.56mg %

Vitamin D status of the study population

- Deficiency < 20ng/ml =167 (53.35%)
- Insufficiency 20 – 30 ng/ml =61 (19.48%)
- Adequacy > 30 ng/ml 84 (26.83%)

Fasting blood glucose > 110mg/dl; N = 38 (12.14%)

Fasting blood glucose between 70 – 110mg/dl; n =274(87.59)

**Table 2** Age wise distribution of vitamin D levels and diabetes

Age wise distribution	N (%)	Mean vitamin D levels	Vitamin D deficiency/ Insufficiency /adequacy n (%)	DM in vitamin D deficiency plus insufficiency (%)
45 – 50	120 (38.33)	26.63	61(19.48)/28(8.94)/31(9.90)	6 (1.91)
51 – 60	113 (36.10)	26.83	72 (23.08)/16(5.11)/25(7.98)	6 (1.91)
> 60	81 (25.887)	26.86	35 (11.18)/17(5.43)/28(8.94)	11 (3.51)

**Table 3** Correlation between vitamin D deficiency and diabetes

DM	N (%)	Normal n (%)
Vitamin D deficiency/ insufficiency	23 (7.34)	206 (65.49)
Vitamin fasting blood glucose levels ±SD (mg%)	176.02 ± 6.56mg	97.65±7.9
Vitamin D adequate	15 (5.11)	68 (21.72)
Mean fasting blood glucose levels ± SD (mg%)	165.42 ± 7.56	99.65 ± 7.9

**DISCUSSION**

Vitamin D deficiency is reported in epidemic proportion, with a prevalence of 70% - 100% in the general population both in urban setting and across all socioeconomic strata.<sup>1</sup>

Study similar to the current study carried in past have reported very high prevalence of vitamin D deficiency among PMW from western world and India.<sup>6-10</sup>

In a study from Europe carried among women aged over 80 years the prevalence of 25 (OH) Inadequacy was 80.9% and 44.5% when considering cut-offs of 75 and 50 n mol/L, respectively.<sup>6</sup>

Results similar to our study were reported among urban central south Chinese PMW indicating 72. 1% women to be vitamin D deficient [25 (OH) D < 50 n mol/L].<sup>14</sup> Vitamin D deficiency was seen in 83.7% of the subjects (< 20ng/ml) at baseline among Indian PMW as reported by one of 82% recent Indian study.<sup>15</sup>

Similarly about 82% of the study group had varying degrees of low 25 (OH) D levels among Indian PMW who were on insufficient calcium and high in phytate diet.<sup>7</sup>

While comparing women of reproductive age (WR) group with PMW both, WR and in PMW 25 (OH) D deficiency (< 20mg/ml), insufficiency (20-30mg/ml) and replete states (> 30 mg/ml) were seen in 76%, 16.5%, 7.5% vs. 70%, 23% and 7% respectively.<sup>16</sup>

Unlike our study PMW with established osteoporoses recorded serum vitamin D to be deficient in two third of patients.<sup>(17)</sup> Thereby indicating that prevention and early detection of hypovitaminosis D is the key to reduce the incidence of osteoporosis among PMW.

In another study no evidence for a clinically important association between serum 25 (OH) D levels and menopause - related symptoms in PMW was observed. The current aspect was beyond the preview of the current study hence not studied.<sup>10</sup>

Interestingly results of current study pointed that commonest age affected with vitamin D deficiency is 50-60 years followed by 45 – 50 and least affected with vitamin D deficiency was above 60 years.

The results are in accordance with the study of [Asadi M et al \(2013\)](#)<sup>18</sup> wherein they showed a statistically significant positive correlation between vitamin D concentration and age in late postmenopausal period. The possible reasons for this may be increasing medical attention in the form vitamin D prophylaxis along with calcium supplementation being received by advancing age people worldwide.

Currently study failed to establish any statistical correlation between vitamin D and diabetes unlike various previous studies. In women, diabetes risk significantly increases in the lowest 25 (OH) D quintile. In men, 25 (OH) D levels has not been shown associated with diabetes incidence.<sup>19</sup>

High prevalence of vitamin D deficiency in patients with type 2 diabetes mellitus (T2DM) and particularly in patients with T2DM and CVD has been reported.<sup>20</sup>

Healthy adults with higher serum 25 (OH) D levels had significantly lower 60 min, 90 min and 129 min postprandial glucose levels and significantly better insulin sensitivity than those who were vitamin D deficient.<sup>21</sup>

The researchers have also reported that with metformin, which improves insulin sensitivity by 13%, higher vitamin D status correlated with a 60% improvement in insulin sensitivity. In a recent clinical trial in women with T2D, vitamin D supplementation was shown to improve insulin sensitivity by 21%.<sup>22</sup>

Another study suggested that vitamin D deficiency is a potential risk factor for obesity and development of insulin resistance leading to T2D.<sup>23</sup>

Most recently [Hiran V et al.](#)<sup>24</sup>, documented that low levels of 25 (OH)D and active 1, 25-Dihydroxyvitamin D is independently associated with T2DM in older Australian men. These variations might be due to the fact that the above mentioned studies were largely epidemiological in nature with different study design taking all men and women without definite age-group into account unlike our study being restricted to PMW.

However, in the study by [Song BM et al.](#)<sup>25</sup>, vitamin D has not been shown to be independently associated with insulin resistance in Korean men and women like our study.

Similarly in a recent meta- analysis of prospective cohort studies no association been between vitamin D intake and T2D has been established.<sup>26</sup>

In another study no correlation was established between 25 (OH) D level with markers of insulin resistance (1R) in postmenopausal Indian women.<sup>15</sup>

The insulin resistance of the obese subjects who were vitamin D deficient and insufficient did not statistically differ from those with vitamin D sufficiency in the study of [Torun E et al.](#)<sup>27</sup>

Low 25 (OH) D levels were not related with higher insulin resistance in obese children and adolescents. In obese subjects, insulin resistance was affected more from BMI.

[Chacko SA et al.](#)<sup>28</sup> documented that higher, serum 25 (OH) D concentrations may be inversely associated with adiposity, triglycerides, triglyceride:high density lipoproteins (HDL) cholesterol ratio, and metabolic syndrome but are not associated with low- density lipoproteins (LDL) and HDL cholesterol, insulin, glucose in PMW.

[Dong JY et al.](#)<sup>29</sup> reported that vitamin D intake during early life although may be associated with a reduced risk of T1D. However, there is not enough evidence for an association between maternal intake of vitamin D and risk of T1D in the off spring.

Sample size of the study group is very small to generalize results of this study group to Indian population. The study has not taken other variable and risk factor into consideration for finding any correlation. However, it addresses a major nutrition related health problem in Indian population, which needs attention.

## CONCLUSION

High prevalence of vitamin D deficiency exists among PMW. However the current study failed to show any statistical correlation between vitamin D deficiency and existence of diabetes, which may be due to small sample size.

## References

1. G R, Gupta A. Vitamin D deficiency in India: Prevalence casualties and inter ventions. *Nutrients* 2014; 6: 729-75.
2. Harinarayan CV, Joshi SR. Vitamin D status in India – Its implications and remedial measures. *J Assoc Physicians India* 2009; 57: 40-8.
3. Marwaha RK, Sripathy G. Vitamin D and bone mineral density of healthy school children in northern India. *Indian J Med Res* 2008; 127: 239-44.
4. Khadilkar AV. Vitamin D deficiency in Indian adolescents. *Indian Paediatr* 2010; 47:756 - 7
5. Sahu M, Bhatia V, Aggarwal A, Rawat V, Saxena P, Pandey A, et al. Vitamin D deficiency in rural girls and pregnant women despite abundant sunshine in northern India. *Clin Endocrinol (Oxf)* 2009; 70:680-4.
6. Bruyere O, sloomian J, Beaudast C, Buckuix, F, cavalier E, Gillain S, et al. prevalence of vitamin D in adequacy in European women aged over 80 years. *Arch gerontol* 2014; 59:78-82.
7. Harinarayan CV. Prevalence of vitamin D insufficiency in postmenopausal south Indian women. *Osteoporosis Int* 2005; 16: 397-402.
8. Paul TV, Thomas N, Seshadri MS, Oommen R, Jose A, Mahendri NV. Prevalence of osteoporosis in ambulatory postmenopausal women from a semi urban region in southern India: A relationship to calcium nutrition and vitamin D status. *Endocr Pract* 2008; 14: 665-71.

9. Kalra S, Kalra B, Sachin Kumar, Khandelwal SK. Vitamin D deficiency in health postmenopausal women in Haryana world *J life Sci Med Res* 2011;1: 11-5.
10. Leblance ES, Desai M, Perrin N, Wactawski-wende J, Manson JE, Cauley JA, *et al.* Vitamin D levels and menopause-related symptoms. *Menopause.* 2014 Nov; 21(11): 1197-1203.
11. Griz LH, Bandeira F, GAbbay MA, Dib SA, Carvalho EF. Vitamin D and diabetes mellitus: An update 2013. *Arq Bras Endocrinol Metabol* 2014; 58:1-8.
12. Dawson- Hughes B, Heaney RP, Holick MF, Lips P, Meunier PJ, Vieth R. Estimates of optimal vitamin D status. *Osteoporosis Int* 2005; 16: 713-6.
13. Hollis BW, Circulating 25- hydroxyl vitamin D levels indicative of vitamin D sufficiency. Implications for establishing a new effective dietary intake recommendation for vitamin D. *J Nutr* 2005; 135:317-22.
14. Lis, ou Y, Zhang H, Zhang Z, Zhou H, Lui L *et al.* Vitamin D status and its relationship with body composition, bone mineral density and fracture risk in urban central south Chinese postmenopausal women. *Ann Nutr Metab* 2014; 64:13-9.
15. Agarwal N, Mithal A, Dhingra V, Kaur P, Godbole MM, Shukhla M. Effect of two different doses of oral cholecalciferol supplementation on serum 25-hydroxy-vitamin D levels in healthy Indian postmenopausal women: A randomized controlled trial, *Indian J Endocrinol Metab* 2013; 17:883-9.
16. Harinaryan CV, Sachan A, Reddy PA, Satish KM, Prasad UV, srivani P, Vitamin D status and bone mineral density in women of reproductive and postmenopausal age groups: A cross-sectional study from south India. *J Assoc physicians India* 2011; 59: 698-704.
17. Nasula R, Tauseef M, Ahmad IA, Agarwal K, Ashok A, Anjana A. Vitamin D deficiency among postmenopausal women with osteoporosis. *J Clin Diag Res* 2013; 7: 336-8.
18. Asadi M, Jouy andeh Z, Nayebzadeh F, Qorbani M. Does aging increase vitamin D serum level in health postmenopausal women? *Acta Med Iran* 2013; 51: 701-4.
19. Schottker B1, Herder C, Rothenbacher D, Perna L, Muller H, Brenner H. Serum 25-hydroxy vitamin D levels and incident diabetes mellitus type 2: A competing risk analysis in a large population based cohort of older adults. *Eur J. Epidemiol* 2013; 28: 267-75.
20. Kavori S, Vuksanove M, Bozovi D, Jovanove M, Jeremi V< Radofici Z, *et al.* Body weight and waist circumference as predictors of vitamin D deficiency in patients with type 2 diabetes and cardiovascular disease. *Vojnosanit Pregl* 2013; 70: 163-9.
21. Chiu KC, Chu A, Go V, Saad MF. Hypovitaminosis D is associated with insulin resistance and beta cell dysfunction. *Am J Clin Nutr* 2004; 79: 820-5.
22. Borissona AM, Tankona T, Kirilov G, Dakovska L, Kovacheva R. The effect of vitamin D3 on insulin secretion peripheral insulin sensitivity in type 2 diabetic patients. *Int J Clin Pract* 2003; 57:258-61.
23. Grineva EN, karonova T, Miceeva E, Belyaeva O, Nikitina 1L. Vitamin D deficiency is a risk factor for obesity and diabetes type 2 in women at late reproductive age. *Aging (Albany NY).* 2013;5:575-81.
24. Hirani V, C umming RG, Le Couteur DG, Naganathan V, Blyth F, Handelsman DJ, *et al.* Low levels of 25-hydroxy Vitamin D and Active 1,25-Dihydroxy vitamin D. Independently associated with type 2 diabetes mellitus in older Australian men: The concord health and ageing in men project. *J Am Geriatr Soc* 2014; Vol. 62, Issue 9: Pages 1741-1747.
25. Song Bm, Kim HC, Choi Dp, Oh SM, Suh 1 Association between serum 25- hydroxyl- Vitamin D levels and insulin resistance in a rural population. *Yonser Med J* 2014; 55: 1036-41.
26. Zhao LM, Tian XQ, Ge JP, Xu YC. Vitamin D intake and type 2 diabetes risk: A meta-analysis of prospective cohort studies. *Afr Health Sci* 2013; 13; 13:1130-8.
27. Torun E, gonullu E, Ozgen 1T, cindemir E, Oktem F. Vitamin D deficiency and insufficiency in obese children and adolescents and its relationship with insulin resistance. *Int J Endocrinol* 2013: 631845.
28. Chacko SA, Song Y, Manson JE, Van Horn L, Eaton C, Martin LW, *et al.* Serum 25-hydroxy vitamin D concentrations in relation to cardiometabolic risk factors and metabolic syndrome in postmenopausal women. *Am J Clin Nutr* 2011; 94: 209-17.
29. Dong JY, Zhang WG, Chen JJ, Zhang ZL, Han SF, Qin LQ. Vitamin D intake and risk of type 1 diabetes: A meta - analysis of observational studies. *Nutrients* 2013; 5: 3551-62.

\*\*\*\*\*

**How to cite this article:**

Shylla Mir and Samiya Mufti.2017, Prevalence of Vitamin D Deficiency among Menopausal Women and Its Correlation with Diabetes: A Cross Sectional Data. *Int J Recent Sci Res.* 8(3), pp. 15909-15912.