



ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

CODEN: IJRSFP (USA)

International Journal of Recent Scientific Research
Vol. 8, Issue, 7, pp. 18635-18643, July, 2017

**International Journal of
Recent Scientific
Research**

DOI: 10.24327/IJRSR

Research Article

A COMPARATIVE ANALYSIS OF PHYSICOCHEMICAL PARAMETERS OF VAIGAI RESERVOIR AND PICKUP DAM, THENI DISTRICT, TAMIL NADU, INDIA

Thamarai Selvi V. P^{1*} and Nirmala T²

Research Centre of Zoology Jayaraj Annapackiam College for Women, Periyakulam

DOI: <http://dx.doi.org/10.24327/ijrsr.2017.0807.0542>

ARTICLE INFO

Article History:

Received 15th April, 2017
Received in revised form 25th
May, 2017
Accepted 28th June, 2017
Published online 28th July, 2017

ABSTRACT

The present study aims at investigating the physicochemical analysis of water samples from two different sites during March - August, 2016 at Vaigai Reservoir and Pickup Dam in Theni District. Evaluation of physicochemical parameters was carried out to assess the quality of water. Each parameter was compared with the standard desirable limit prescribed by Bureau of Indian Standards (BIS). The physicochemical parameters of Vaigai Reservoir and Pickup Dam water sample was well within the limit where as Manganese, Ammonia, Nitrite and Phosphate parameters were not within the permissible limit.

Key Words:

Physicochemical parameters, Vaigai Reservoir, Pickup Dam and water quality.

Copyright © Thamarai Selvi V. P and Nirmala T, 2017, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Water is one of the most important and abundant compounds of the ecosystem. It is also a crucial resource for agriculture, manufacturing sites and other human activities. It is the most essential basic component to all living beings. As most of the biochemical reactions that takes place through the metabolism and growth of living organisms required water, a vital resource required for their growth. River is the most important and vulnerable freshwater system that plays a critical role in the sustenance of all life. In urban areas, the careless disposal of industrial effluents and other wastes in rivers and lakes may contribute greatly to the poor quality of river water (Agarwal and Sexana, 2011).

Water is considered as one of the nutrients, although it yields no calories, yet it enters into structural composition of the cell and is an essential component of diet (Baloch *et al.*, 2000). WHO (2001) reported that approximately 36% of urban and 65% of rural Indians are without access to safe drinking water (Akoto and Adiyah, 2007). Fresh water is one of the most important resources for the survival of all the living organisms. It is even more important for the human beings as they depend upon it for food production, industrial and waste disposal as well as cultural requirement (Akpoveta *et al.*, 2011). It is estimated that over 70 per cent of India's food grain production comes from irrigated agriculture, in which ground water plays a

major role (Gandhi and Namboodiri, 2009). In India, 6.73 hectare of land is affected by salinity and sodicity (Singh *et al.*, 2009). Groundwater contains a varying amount of different kinds of ions such as carbonate, bicarbonate, calcium, magnesium, sulphate and hardness (Choudhary and Dagankar, 2007).

The need for water quality could be determined in different ways either for drinking, other domestic uses, industrial, agricultural, irrigation or fish farming. However, the maintenance of aquatic ecosystem is totally dependent on the physicochemical properties and biological diversity (Verkatesharajlu *et al.*, 2010). The level of physicochemical parameters of water will determine the purpose with which the water could be best used for with little or no treatments. Variability in physicochemical parameters is responsible for the distribution of organisms in different freshwater habitat according to their adaptation, which allow them to survive in a specific habitat (Jeffries and Mills, 1990). Major shift in the stream bed composition and processes can alter species distribution productivity (Palmer *et al.*, 1997).

The search for good portable water has been man's utmost concern since the beginning of civilization and thus, scarcity of clean freshwater is one of the world most pressing environmental problems. Therefore, in search of domestic water which includes the water needs for families for drinking,

*Corresponding author: Thamarai Selvi V. P

Research Centre of Zoology Jayaraj Annapackiam College for Women, Periyakulam

cooking, sanitation, hygiene etc, (Bustanmante et al., 2004), rural community dwellers in the river front depend on the rivers or streams as a source of water for their domestic and other uses. This could be as a result of uneasy access and high cost of harnessing underground water.

Good quality of water resources depends on a large number of physico chemical and biological characteristics. These parameters are essential to identify magnitude and source of any pollution load (Ushe and Amadi, 2005).

The protection of ground water from surface activity is very difficult to remediate but risk can be minimized by monitoring of highly pollution prone area. Monitoring program is reliable for estimation of water quality (Chowdhury et al., 2017). Hence this study has been made to analyse the Physico chemical parameters of water samples from Vaigai Reservoir of Theni Districts.

MATERIALS AND METHODS

Description of the Study Site

The study has been carried out in two sampling sites such as Vaigai Reservoir and Pickup Dam in Theni District, Tamil Nadu during March - August 2016.

Vaigai Reservoir originates in the Periyar Plateau of the Western Ghats range, and flows to Northeast through the Cambam Valley, which lies between the Palni Hills to the north and Varushanad Hills to the South. The Vaigai is 258 kilometers (160m) long, with a drainage basin of 7,031 square kilometers (2,715 sq m) large. It falls within the coordinates of Latitude at 7°21'00 N and Longitude 79°00'00 E. Vaigai Dam is built across the Vaigai Reservoir near Andipatty, in Theni district of Tamil Nadu, Southern India. The Vaigai Reservoir rarely floods. Its chief tributaries are the Suruliar, Theniar, Varaha Nadhi and it supplies water to Madurai, Sivaganga and Ramanathapuram districts.



Plate 1 Vaigai Reservoir



Plate 2 Pickup Dam

Sample Collection and Preservation

Water samples were collected in sterilized plastic bottles for every month in the early hours during March - August, 2016.

Analysis of Water Samples

Colour, Odour, pH and Temperature of the water samples were measured on the sampling spot by water analyzer kit. Total alkalinity, total hardness, magnesium, calcium and chloride contents were analyzed by titrimetric methods in the laboratory following the standard methods as prescribed by APHA, (2005).

RESULTS AND DISCUSSION

The sample was slightly turbid, greenish and yellow colour was observed with a fowl smell. Water temperature ranged from 20 - 32°C, air temperature varied between 23 and 34°C. Highest temperature in both air and water was recorded in Pickup Dam during July. Rainfall was observed during July low in August. There is no rainfall in other months (Table 1).

Physical Parameter in Water samples from Vaigai Reservoir and Pickup Dam

Turbidity

Turbidity was high in June and it was low in March. In both the sites, high turbidity was observed in June, July and August and it was low in May. Vaigai Reservoir showed high turbidity in four months except April and May (Figure 1) when compared to Pickup Dam Similar results were also observed by Manjare et al. (2010).

Water turbidity, which reflects transparency, is an important criterion for assessing the quality of water. This might be due to improper disposal of sewage, surface runoff and wastewater from different domestic activities. Similar higher turbidity values are also recorded by many workers as compared to the limit set by WHO (Akan et al., 2008; Joshi et al., 2009; Mebrahtu and Zerabruk, 2011; Murhekar, 2011; Pal et al., 2013).




Figure 1 Turbidity of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Total Dissolved Solids

Total dissolved solids indicate the salinity of groundwater. High value of total dissolved solid was observed in June and low in March in both the sampling sites. To compare with high total dissolved solid was observed in Pickup Dam (Figure 2).

Table 1 Physical Parameters in Water samples from Vaigai Reservoir and Pickup Dam during March-August, 2016

Month	Site	Appearance	Colour	Odour	Air Temp (°C)	Water Temp (°C)	Rainfall (Mm)
March	Reservoir	Slightly Turbid	Greenish in colour	Slight Fowl Smell	23	20	0
	Pickup Dam	slightly Turbid	Clear and Colourless	Fowl Smell	26	24	0
April	Reservoir	Slightly Turbid	Greenish	Fowl Smell	24	22	0
	Pickup Dam	slightly Turbid	Greenish in colour	Fowl Smell	27	25	0
May	Reservoir	Slightly Turbid	Greenish in colour	Fowl Smell	26	23	0
	Pickup Dam	slightly Turbid	Greenish in colour	Fowl Smell	29	27	0
June	Reservoir	Slightly Turbid	Greenish in colour	Fowl Smell	27	22	0
	Pickup Dam	slightly Turbid	Greenish Yellow	Slight Fowl smell	34	32	0
July	Reservoir	Slightly Turbid	Yellow in colour	Fowl Smell	26	20	59
	Pickup Dam	slightly Turbid	Clear and Colourless	Fowl Smell	37	29	59
August	Reservoir	Slightly Turbid	Greenish Yellow	Fowl Smell	26	20	4
	Pickup Dam	slightly Turbid	Greenish Yellow	Slightly Fowl smell	34	32	4

Total dissolved solids in two samples were below the permissible limit, whereas in the study of [Kanase et al. \(2016\)](#) showed higher values of total dissolved solids. The value of the present study agrees with the result of [Giorgis et al. \(2010\)](#) and [Alemayehu \(2008\)](#). In Dhamola River total dissolved solids were within the permissible limit in all the sampling sites ([Dinkar, 2016](#)). Water containing more than 500 mg/l of total dissolved solid is not considered desirable for drinking water supplies, but in unavoidable cases 1500 mg/l is also allowed ([Shrinivasa and Venkateswaralu 2000](#)).




Figure 2 Total Dissolved Solids of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Electrical Conductivity [EC]

Electrical Conductivity was high in June and low in March. When combine the two sites the Pickup Dam showed high electrical conductivity and Vaigai Reservoir showed low electrical conductivity in all the months. Hence EC was high in Pickup Dam than in Vaigai Reservoir (Figure 3). [Jayalakshmi et al. \(2011\)](#) reported that electrical conductivity values of Elala River were within standard permissible unit which are similar to our present findings. Increase in electrical conductivity indicates the presence of higher concentration of ions ([Deepali et al., 2009](#)).

The higher levels of electrical conductivity alter the chelating properties of receiving systems, which favour free metal availability to flora and fauna ([Nagajyothi et al., 2009](#)).




Figure 3 Electrical Conductivity of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

[Navneeth Kumar et al, \(2010\)](#) suggested that the underground drinking water quality of study area can be checked effectively by controlling conductivity of water and this may also be applied to the management of water quality.

p^H

pH of the Vaigai Reservoir water samples was high in April and it was low in March. In Pickup Dam the pH value was high in May and low in June. Among the two sampling sites, the water sample from the Vaigai Reservoir showed high alkalinity (Figure 4). Similar observations made by [Venkateswarlu et al. \(2014\)](#), [Tyor and Chawla, \(2012\)](#); and [Chaudhary et al, \(2013\)](#). pH is more important in determining the corrosive nature of water. Lower the pH value higher is the corrosive nature of water. pH was positively correlated with electrical conductance and total alkalinity ([Gupta 2009](#)). The higher pH values observed suggested that carbon dioxide, carbonate-bicarbonate equilibrium is was affected due to change in physico-chemical condition.




Figure 4 pH of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

maximum level of total hardness is due to the presence of carbonate and non carbonate compounds (Ramesh *et al.*, 2013).




Figure 6 Total Hardness of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Total Alkalinity

Alkalinity was high in June and low in March in both the sites (Figure 5). High alkalinity was observed by Venkateswarlu *et al.* (2014); Alemayehu (2001); Sharma *et al.* (2013); Soni *et al.* (2013). Lower values of alkalinity was reported by Bhattarai *et al.* (2008); Joshi *et al.* (2009); Alani *et al.* (2014). The alkalinity of both the water samples were within the standard permissible unit of BIS.




Figure 5 Total Alkalinity of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Calcium

Calcium was high in May and it was low in March in both the sampling sites. Comparatively high calcium was present in Pickup Dam (Figure 7) the present study coincides with the study of Vanitha and Shunmugavelu, (2012) and Kumar *et al.* (2004).

Calcium is directly related to hardness. Calcium may dissolve readily from carbonate rocks and lime stones (Chari and Lavanya, 1994). Calcium concentration ranged between 12 mg/l to 172 mg/l and found below the permissible limit of ISI. Calcium ions usually increase the hardness content in the river water (Prasad, 2016).




Figure 7 Calcium of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Total Hardness

Total hardness was high in April, June, July and August and it was low in March in Vaigai Reservoir. In Pickup Dam it was high in June and April and was low in March, July and August (Figure 6). The standard permissible unit of values was observed by Prasad *et al.* (2016). They point out the concordant values of hardness. Mishra *et al.* (2014) studied the pond water quality in Rairangpur, Varanasi and found that the hardness values varied from 146 to 268mg/l.

Hardness of water is a measure of its capacity to form precipitates with soap and scales with certain anions present in the water. Water hardness arises due to the presence of cations such as calcium, magnesium and anions such as bicarbonates, chlorides and sulfides (Kumar *et al.*, 2013; Adejuwon and Adelakun, 2012). Water hardness is understood as a measure of the capacity of water to precipitate soap. The increase in the

Magnesium

In Vaigai Reservoir, magnesium content was high in April, June, July and August and it was low in March and May. When comparing the two sampling sites, magnesium was observed high in Pickup Dam than Vaigai Reservoir. The magnesium content was within the permissible limit in both the sampling sites (Figure 8). The reports of Balamurugan (2015) and Chittora *et al.* (2017) do not fall in line with the present investigation. Vanitha (2012) has reported that the results, which reveal that magnesium concentration, lie within the permissible limit at Kunnur and Vaigai dam Sample units. Magnesium is one of the important sources for chlorophyll

development and it acts as a restrictive factor for the growth of phytoplankton (Dagaonkar and Saksena, 1992).




Figure 8 Magnesium of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Ammonia

Ammonia was high in June and it was low in May, July and August in both the sampling sites. To compare the two samples, high ammonia was observed in Pickup Dam than Vaigai Reservoir (Figure 9). The ammonia was above the desirable limits of BIS standard. Similar result was reported by Seyoum, (2007). Ammonia is frequently present in groundwater sources where there is no oxygen present. Ammonia ions play a key part in water treatment because they need to be removed before breakpoint chlorination can be achieved. Chittora *et al.* (2017) reported 0.03-150mg/l of ammonia in the drinking water sample Udaipur.




Figure 9 Ammonia of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Nitrite

Nitrite was high in June and low in May, July and August in Reservoir. In Pickup Dam high nitrate content was observed in June and it was low April, May, July and August (Figure 10) similar results were observed by Drissi (2016); Sivamanikandan and Ahmed John, (2016). Nitrite exists normally in very low concentration and even in waste treatment plant, effluent levels are relatively low, principally because the nitrogen will tend to exist in the more reduced (ammonia) or more oxidized (nitrate) forms. Nitrite is an intermediate in the oxidation of ammonia to nitrate, such oxidation can proceed in soil, and because sewage is a rich source of ammonia, waters which show any appreciable amounts of nitrite are regarded highly poor in quality. Levels in unpolluted waters are

normally low, that is below 0.03 mg/dm³ NO₂. Values greater than this indicate sewage pollution WHO 1993.




Figure 10 Nitrate of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Nitrate

Nitrate content was high in May and July and it was low in March, April June and August in both the sampling sites (Figure 11). Concordant values were recorded by Alam (2017). Nitrate represents the final product of biochemical oxidation of ammonia (Mahananda *et al.*, 2010). Similar results were recorded in Sivamanikandan and Ahmed John, (2016). Nitrate is the most important nutrient in an ecosystem. Generally water bodies polluted by organic matter exhibit higher values of nitrate. Moreover nitrogen is an essential nutrient that is required by all plants and animals for the formation of amino acids. In its molecular form, nitrogen cannot be used by most aquatic plants, therefore it must be converted to another form. One such form is ammonia.




Figure 11 Nitrate of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Chloride

Chloride content was high in June and it was low in on other months in the Vaigai Reservoir. In Pickup Dam it was high in June was low in March April July and August (Figure 12). Permissible limit of chloride according to BIS is 250 -1000 mg/l and it is within the limit. Similar results were reported by Swarnalatha and Narsingrao, (1998) and Manjare *et al.* (2010) where the higher value was observed by Mallika (2017).

Chloride is one of the common anions found in water and sewage. Chloride is highly soluble with most of the naturally occurring cations and do not precipitate with sediments and cannot be removed biologically in the treatment of waste. The

concentration of Chloride in natural waters varies from a few milligrams to several thousand milligrams per liter.




Figure 12 Chloride of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Fluoride

Fluoride was high in June and low in May in the Vaigai Reservoir, whereas in Pickup Dam, it was high in April and June and it was low in March, July and August (Figure 13). The concentration of fluoride was within the standard permissible limit as in the study of Kumar and Kumar (2013); Kumar and Jain (2016). Fluoride occurs as fluorspar (fluorite), rock phosphate, triphite, phosphorite crystals etc. in nature. Among factors which control the concentration of fluoride are the climate of the area and the presence of accessory minerals in the rock mineral assemblage through which the ground water is circulating (Handa, 1985).




Figure 13 Fluoride of water samples from Vaigai Reservoir and Pickup dam during March-July, 2016

Sulphate

The presence of sulphate concentration was maximum in May and it was minimum in March and June in the Vaigai Reservoir. In Pickup Dam, it was high in June and low in except all the months (Figure 14). The sulphate content was below the permissible limit. The concordant values were observed by Lakshmi, (2017) in the sulphate content.

The natural concentration of sulphate in most surface water is within the range of 2 to 80 mg/l (Manivasakam, 2005). Sulphate is widely distributed in nature and may be present in natural waters. The main source of sulphur is the rocks present near the water bodies and biochemical action of anaerobic bacteria (Sharma et al., 2011). Anbarasu and Anbuselvan,

(2017) reported by higher values of sulphate content in Musiri Taluk unlike this study area.




Figure 14 Sulphate of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Phosphate

Phosphate was high in June and low in April and May in Vaigai Reservoir. In Pickup Dam, it was high in March and low in April and March (Figure 15). Phosphate content was present in above the permissible limit. This finding is in agreement with that of Udaipur lakes (Ranu 2001, Christy 2002 and Medudhula et al., 2012). It is one of the most important nutrients and a limiting factor in the maintenance of reservoir fertility. Phosphate may occur in ground water as a result of domestic sewage, detergents, agricultural with fertilizers and industrial waste water. The phosphate content in the Vuyyuru, Andra Pradesh was found in the range of 0.31 mg/l to 0.66 mg/l and it was within the limit (Janardhana et al., 2013).




Figure 15 Phosphate of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

Dissolved Oxygen

Dissolved oxygen was high in June and low in May in the Vaigai Reservoir. In Pickup Dam also it was high in June and low in March, July and August (Figure 16). Similar results of dissolved oxygen were observed in Medudhula, (2012). Dissolved Oxygen is one of the most important parameter. Its correlation with water body gives direct and indirect information e.g. bacterial activity, photosynthesis, availability of nutrients, stratification etc. (Vikal, 2009). Dissolved oxygen is an important aquatic parameter, whose presence is vital to aquatic fauna. It plays crucial role in life processes of animals. Rani et al, (2004) also reported lower values of dissolved oxygen in summer season due to higher rate of decomposition

of organic matter and limited flow of water in low holding environment due to high temperature.




Figure 16 Dissolved oxygen of water samples from Vaigai Reservoir and Pickup dam during March-August, 2016

CONCLUSION

The major intention of physicochemical analysis is to prevent physical changes as well as chemical contamination of reservoir so as to protect the life of people in our society. In this study, two samples were collected from two different locations of Theni District. The physicochemical analysis of Vaigai Reservoir and Pickup Dam water was analysed by desirable, permissible and acceptance limit of the water sample. The parameters of turbidity, pH, electrical conductivity and fluoride are within the desirable and permissible limits. Total alkalinity, total hardness, calcium, magnesium and nitrite are below the permissible limit, phosphate, ammonia and chloride are above the permissible and desirable limit. Therefore, it is recommended in this study, to create awareness among the residents of the studied area about the need to utilize Vaigai Reservoir and Pickup Dam water for irrigation and domestic purposes.

References

Abbhijit Chowdhury, Pulak Kumar Patra and and Shibani Chaudhury 2017. Physico-chemical characterization of water quality in the Bastacola coalmine regions of Jharia coalfield, Dhanbad, India, *Int. Res. J. Earth Sci*, Vol. 5(2), 1-8.

Adejuwon J. O., and Adelakun M. A., 2012. Physiochemical and bacteriological analysis of surface water in Ewekoro Local Government Area of Ogun State, Nigeria: Case study of Lala, Yobo and Agodo Rivers, *International Journal of Water Resources and Environmental Engineering*, 4(3), 66-72

Agarwal A and Saxena 2011. M Assessment of pollution by physicochemical water parameters. *Advanced Applied Science Research* 2(2) 185-189.

Akan J.C, Ogugbaja V.O, Abdulrahmen F.I, Ayodele J.T 2008. Pollution Levels in Effluent Samples from Tanneries and Textile of Kano Industrial Areas, Nigeria; *Global Journal of Pure and Applied Sciences* Vol 15, No. 3 PP 343-352.

Akan JC, Ogugbaja VO, Abdulrahmen FI, Ayodele JT 2008. Pollution Levels in Effluent Samples from Tanneries and Textile of Kano Industrial Areas, Nigeria; *Global Journal of Pure and Applied Sciences* Vol 15, No. 3 PP 343-352.

Akoto, O and Adiyiah, J. 2007. Chemical analysis of drinking water from some communities in the brong ahafo region.

International Journal of Environmental Science and Technology, 4(2):211-214.

Akpoveta, O.V., F. Egharevba, O.W. Medjor, O.K. Ize-Iyamu, and E.D. Enyemike. 2011. "Microbial Degradation and its Kinetics on Crude Oil Polluted Soil". *Research Journal of Chemical Sciences*. 1(6):8-14

Alani, R., Alo, B and Ukoakonam, F. 2014. Preliminary Investigation of the State of Pollution of Ogun River at Kara Abattoir, Near Berger, Lagos. *International Journal of Environmental Science and Toxicology*, 2(2): 11-23.

Alemayehu, T. 2001. The impact of uncontrolled waste disposal on surface water quality in Addis Ababa, Ethiopia. *SINET Ethiopian Journal of Science*, 24(1), 93-104.

Alemayehu, T. 2008. Ethiopia spate irrigation country profile. Addis Ababa, Oromia Water Works Supervision and Design Enterprise

Amadi, A.N., P.I. Olasehinde and J. Yisa, 2010. Characterization of Groundwater chemistry in the Coastal plain-sand Aquifer of Owerri using Factor Analysis. *Int. J. Phys. Sci.*, 5: 1306-1314.

APHA 2005. Standard methods for the examination of water and waste water. 21st ed. American Public Health Association, Washington, DC, USA.

Balamurugan, Sivakumar. The seasonal variation of water quality parameters in Vaigai River at Madurai. *Int.Journ. Comp. and Org. Trends*. 2015; 19(1):42-50.

Baloch M.K, Jan IU and Ashour S.T 2000. Effect of septic tank effluents on quality of ground water. *Pakistan Journal of Food Sciences*, 10(3-4):25-31.

Bhattarai, K. R., Shrestha, B. B and Lekhak, H. D. 2008. Water Quality of Sundarijal Reservoir and its Feeding Streams in Kathmandu. *Scientific World*, 6(6): 99-106.

Bustanmante R, Butterworth J, Flierman M, Herbas D, Hollander M, Meer S.V, Ravenstijn P, Reynanga M and Zurita.G 2004. Livelihoods in conflict dispute over water for household level productivity uses in Tarata Blivia. *Intl.water and sanit. publ., Delf Netherlands*.

Chari K.V.R. and Lavanya M.G., 1994. Groundwater contamination in Cuddapah urban area, Andhra Pradesh, In Proceedings on regional Workshop of Environmental aspects of groundwater development. KU, Kurukshetra Oct. 17-19, Kurukshetra, India, 130-134.

Chaudhari, P.R., Ahire, D.V., Ahire, V.D., Chkravarty, M. and Maity, S. 2013. Soil Bulk Density as related to Soil Texture, Organic Matter Content and available total Nutrients of Coimbatore Soil. *Int. J. Scientific and Res. Publications*. 3(2)

Chisty. N., 2002. Studies on Biodiversity of Freshwater Zooplankton in Relation to Toxicity of selected Heavy Metals. Ph. D. Thesis submitted to M.L Sukhadia Univeristy Udaipur.

Chittora A.K., 2017Chandra Shekar Kapoorand Vidhya Kapasya, Comparative Assessment of Physicochemical Parameters of Udaipur City, (Raj.) India, *Int J Environ Sci Nat Res*, 1(2).

Choudhary P, DagankarA and Praveen S, 2007. Physicochemical analysis of groundwater for evaluation of drinking water quality at Dhar, M.P. *National environmental and Pollution Technology, Technoscience Pub*. 6(1):109-112.

- Dagaonkar and D.N. Saksena, 1992. Physicochemical and biological characterization of a temple tank, Kaila Sagar, Gwalior, Madhya Pradesh. *J. Hydrobiol.* 8 (1), 11-19.
- Dhanaji Kanase G, Shagufta Shaikh A, Pramod Jagadale N 2016. Physico-Chemical Analysis of Drinking Water Samples of Different Places in Kadegaon Tahsil, Maharashtra (India), Pelagia Research Library Advances in Applied Science Research, 7(6):41-44.
- Ermias Seyoum, 2007. Assessment of the Physicochemical Parameters of River Beressa in Debre Berhan Town (North Shoa) for suitability of drinking water, M. Sc. Thesis
- G. Vanitha. M. Shunmugavelu 2012. Hydrochemistry and Environmental Status of Vaigai River Water in Tamil Nadu, India, *International Journal of Engineering Research and Applications* Vol. 2, Issue 5, pp. 1942-1946.
- Gandhi V.P and Namboodiri N.V. Groundwater Irrigation in India: Gains, Costs and Risks. Indian Institute of Management, Ahmedabad, India. 2009.
- Giorgis, Y.H., Lemma, B., Birhanu, B., Sahu, K. and Sintayehu, W. 2010. Environmental impact assessment and policy on Lake Hawassa, SNNPRs. SOS-Sahel, Ethiopia. Draft document.
- Gupta, D. P., Sunita and J. P. Saharan 2009. Physicochemical Analysis of Ground Water of Selected Area of Kaithal City (Haryana) India, *Researcher*, 1(2), pp 1-5.
- Handa, A.K, Pasrija, H.B.D, Shrivastava, A.K, Jain Neeraj, 1997. Ecological studies on Sukhana Lake. *Env. Eco*, 15(2), 370-373.
- Janardhana Rao D., Hari Babu B, Swami A.V.V.S and Sumithra S, 2013. Physico-Chemical Characteristics of Ground Water of Vuyyuru, Part of East Coast of India, *Universal Journal of Environmental Research and Technology*, 3(2): 225-232.
- Jayalakshmi, V., Lakshmi, N and Singara Charya, M. A. 2011. Assessment of physicochemical parameters of water and waste waters in and around Vijayawada. *International journal of research Pharmaceutical and Biomedical Sciences*, 2(3): 1040-1046.
- Jayalakshmi, V., Lakshmi, N and Singara Charya, M. A. 2011. Assessment of physicochemical parameters of water and waste waters in and around Vijayawada. *International journal of research Pharmaceutical and Biomedical Sciences*, 2(3): 1040-1046.
- Jeffries, M. and Mills, D., 1990. Freshwater Ecology. Principles and Applications. pp: 335-337. Belhaven Press, London and New York.
- Joshi V.J. and Santani D.D.2012. Physicochemical Characterization and Heavy Metal Concentration in Effluent of Textile Industry, *Universal Journal of Environmental Research and Technology*, 2(2), 93-96.
- K. Anbarasu, G. Anbuselvan 2017. Physico-chemical parameter analysis of water in Musiri Taluk, Tamil Nadu, India. *World News of Natural Sciences* 6: 36-43
- K. Hari Prasad, C. Anjali, M. Sridhar Reddy 2016. Study of Physico-Chemical Characteristics of Penna River at Chennai, Cuddapah Basin, India, *Indian Journal of Advances in Chemical Science* 4(1) 20-24.
- K. Hari Prasad, C. Anjali, M. Sridhar Reddy, 2016. Study of Physico-Chemical Characteristics of Penna River at Chennai, Cuddapah Basin, India, *Indian Journal of Advances in Chemical Science* 4(1) 20-24.
- Kumar M, Kumar R 2014. Assessment of physico-chemical properties of ground water in granite mining area in Goramachia, Jhansi (India). *Int Res J Environment Sci* 2: 19-24.
- Kumar R., Singh, R.D., Sharma, K.D., 2004, "Water Resources of India", *Current Science*, 89(5), pp 794-811.
- M. F. Alam, N. C. Dafader, S. Sultana, N. Rahman and T. 2017. Taheri, Physico-Chemical Analysis of the Bottled Drinking Water available in the Dhaka City of Bangladesh, *Journal of Materials and Environmental Sciences, JMES*, 8 (6), pp. 2076-2083
- M.R.Mahananda, B.P.Mohanty and N.R. Behera 2010. Physico-Chemical Analysis of Surface and Ground Water of Bargarh District, Orissa, India *JRRAS* 2 (3).
- Manivasakam N., 2005. Physicochemical examination of water sewage and industrial Effluent, 5 th Edition, ISBN: 978-0-8206-0040-6, Pragati Prakashan Meerut, 441p.
- Manjare S. A; Vhanalakar S. A and Muley D. V 2010. Analysis of water quality using physicochemical parameters tamdalge tank in Kolhapur district, Maharashtra. *Int. J. Adv. Biotech.Res.* 1(2): 115-119.
- Manjare S. A; Vhanalakar S. A. and Muley D. V. 2010. Analysis of water quality using physicochemical parameters tamdalge tank in Kolhapur district, Maharashtra. *Int. J. Adv. Biotech.Res.* 1(2): 115-119
- Mebrahtu G., Zerabruk S., 2011. Concentration and Health Implication of Heavy Metals in Drinking Water from Urban Areas of Tigray Region, Northern Ethiopia. *Momona Ethiopian Journal of Science.* 3(1):105-21.
- Medudhula.Thirupathaiiah, Ch.Samatha, Chintha Sammaiah, 2012. Analysis of water quality using physico-chemical parameters in lower manair reservoir of Karimnagar district, Andhra Pradesh; *Int J OF Env Sciences*, 3(1).
- Murhekar Gopalkrushna H 2011. International Journal of Research in Chemistry and Environment; Murhekar Gopalkrushna *Int. J. Res. Chem. Environ.* 1(2) (183-187).
- Nagajyothi P C; Dinakar N; Suresh S; Udaykiran Y; Suresh C and Damodharan T 2009. Effect of industrial effluent on the morphological parameters and chlorophyll content of green gram (*Phaseolusaureus* Roxb). *J. Environ. Biol.* 30: 385-388
- Navneet Kumar, D. K. Sinha, 2010. Drinking water quality management through correlation studies among various physicochemical parameters: A case study, *International Journal of Environmental Sciences*, 1(2), pp 253-259
- P. Ravi Kumar, M. Aneesul Mehmood, K. K. Somashekar, 2013. Water quality index to determine the surface water quality of Sankey tank and Mallathahalli Lake, Bangalore Urban district, Karnataka, India, *Applied Water Science*, 3: 247-261.
- Palmer M.A, Alan P.C, Bland J.F, Janine G, Kevin D.H, Richard K.J, Timo K, Sam L, Charles R.L, Robert J.N, Claudia R, Francesc S and David S (1997). Biodiversity and eco-system processes in freshwater sediments *Ambio*. 26: 571-577.
- Premlata, Vikal, 2009. Multivariate analysis of drinking water quality parameters of lake Pichhola in Udaipur, India.

- Biological Forum, *Biological Forum- An International Journal*, 1(2), pp 97-102.
- Ramesh M; Dharmaraj E and Raj B. J. R 2013. Physico-chemical characteristics of ground water of Manachanallur Block Trichy, Tamil Nadu, India. *Adv. App. Sci. Res.* 3(3):1709-1713.
- Rani, R., Gupta, B.K and Srivastava, K.B.L., 2004. Studies on water quality assessment in Satna city (M.P): Seasonal parametric variations, *Nature environment and pollution technology*, 3(4), pp 563-565.
- Ranu 2001. Studies on toxicity of textile effluents to fresh water Zooplankton. Ph.D. Thesis submitted to MLSU, Udaipur.
- Rima Drissi and Chaabane Mouats 2016. Physicochemical Analysis of Water of Boumerzoug River (Constantine-Algeria), *Research Journal of Pharmaceutical, Biological and Chemical Sciences* 7(1).
- S Mallika, R Umamaheswari and S Krishnamoorthy 2017. Physico-Chemical parameters and bacteriological study of vaigai River Water Madurai district, Tamilnadu, India, *International Journal of Fisheries and Aquatic Studies*, 5(1): 42-45.
- S. Mishra, A. L. Singh and D. Tiwary 2014. Studies of Physico-chemical Status of the Ponds at Varanasi Holy City under Anthropogenic Influences. *Int. J. Environ. Res. Devel.* 4: 261-268.
- Sharma D. and Kansal A. 2011. Water quality analysis of river Yamuna using water quality index in the national capital territory, India (2000-2009). *Appl. Wat. Sci.*, 1(3-4), 147-157.
- Singh A.K, Qadar A and N.P.S Yadvansh 2009. Mineral nutrition of crop plants in salt affected areas in India. ICAR, New Delhi, CSSRI, Karnal.
- Sivamanikandan, P., and Ahmed John, S. 2016. Impact of physico-chemical parameters on bacterial population in Mullaiperiyar River water-Theni district, Tamilnadu, India. *Afr.J.Microbiol.Res*, 9(1): 26-32.
- Sivamanikandan, P., and Ahmed John, S. 2016. Impact of physico-chemical parameters on bacterial population in Mullaiperiyar River water-Theni district, Tamilnadu, India. *Afr.J.Microbiol.Res*, 9(1): 26-32.
- Soni, H. B., Dabhi, M and Thomas, S. 2013. Surface water quality assessment and conservation measures of two pond ecosystems of Central Gujarat. *International Research Journal of Chemistry*, 3(3): 69-81.
- SrinivasaRao B and Venkateswarlu P. 2000. Evaluation of ground water quality in Chirala town (Prakasam district). *Ind. J. Environ. Protec.*: 20(3):161-164.
- Swaranlatha, S. and A.Narsingrao., (1998), Ecological studies of Banjara lake with reference to water pollution, *Journal of environmental biology*, 19(2), pp 179-186.
- Tyor A.K. and D. Chawla 2012. Survey and study of phytoplankton ecology in Sukhna lake, Chandigarh, India. *Intl J Appl Biol Pharma Tech.* 3(2): 229-237.
- Venkatesharaju K., Ravikumar.P., Somashekar. R.K., Prakash. K.L., 2010, Physico-chemical and Bacteriological Investigation on the river Cauvery of Kollegal Stretch in Karnataka, *Journal of science Engineering and technology*, 6 (1), pp 50-59.
- Venkateswarlu P., K. Chandra Mohan and J. Suresh. 2014. Physio-chemical analysis of bore-well water of Kurnool environs, Andhra, *Journal of Chemical and Pharmaceutical Research*, 6(9):77-80.
- WHO, (1993) World Health Organization. Guidelines for Drinking Water Quality, World Health Organization, Geneva, Switzerland.

How to cite this article:

Thamarai Selvi V. P and Nirmala T., A Comparative Analysis of Physicochemical Parameters of Vaigai Reservoir And Pickup Dam, Theni District, Tamil Nadu, India. *Int J Recent Sci Res.* 8(7), pp. 18635-18643.
DOI: <http://dx.doi.org/10.24327/ijrsr.2017.0807.0542>
