



ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

CODEN: IJRSFP (USA)

International Journal of Recent Scientific Research
Vol. 9, Issue, 1(C), pp. 23083-23086, January, 2018

**International Journal of
Recent Scientific
Research**

DOI: 10.24327/IJRSR

Research Article

DIVERSITY OF ENDOPHYTIC FUNGI FROM LEAVES OF *Centella asiatica* L. and *Melia azedarach* L

Shijila Rani A.S., Ambikapathy, V and Panneerselvam A

Department of Botany and Microbiology, AVVM Sri Pushpam College (Autonomous),
Poondi, 613 503, Thanjavur

DOI: <http://dx.doi.org/10.24327/ijrsr.2018.0901.1394>

ARTICLE INFO

Article History:

Received 17th October, 2017
Received in revised form 21st
November, 2017
Accepted 05th December, 2017
Published online 28th January, 2018

Key Words:

Centella asiatica., *Melia azedarach*.,
leaves, Czapek dextrose agar

ABSTRACT

Endophytic fungi are symbiotic association and the harbor of plants. It is living in inner tissues and without causing any apparent symptoms. In the present investigation, the endophytic fungi were isolated from leaves of *Centella asiatica* and *Melia azedarach*. These two plants are highly medicinal properties. Totally eight different genera were isolated from both the leaves. Generally, *Absidia repens*, *Alternaria* sp., *Aspergillus flavus*, *A. terreus*, *Chaetomium globosum*, *Curvularia* sp., *Fusarium solani*, and *Humicola* sp. The maximum percentage of endophytic fungi and diversity indices were recorded in *Melia azedarach* leaves fragment when compared to *Centella asiatica* leaves.

Copyright © Shijila Rani A.S., Ambikapathy, V and Panneerselvam A, 2018, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Endophytes are ubiquitous in nature and are found in every plant. Endophytic fungi are microorganisms found inside plant species (Stone *et al.*, 2004) and may play a key role in plant development by controlling endophytic microbes and herbivorous insects or producing growth promoting substances (Neto *et al.*, 2002). Endophytic fungi also have functions related to saprophytic microorganisms, and many species are involved in the process of maturation and leaf decomposition (Promputtha *et al.*, 2007; Sieber, 2007). Endophytic fungi infect and inhabit primarily the aerial tissue of the host plant without causing detectable symptoms. The relationship between endophytes are thought to be symbiotic, such as those endophytes to get nutrients and protection from the host but contribute to effective host defense mechanism against pathogens herbivores or abiotic stress (Redman *et al.*, 20012 and Arnold and Herre, 2003).

One of the important role of endophytic fungi is to initiate the biological degradation of dead or dying host plant which is necessary for nutrient recycling (Strobel, 2000). Endophytic fungi are reported from plants grow in various environmental including tropic (Mohali, *et al.*, 2005), temperate (Ganley *et al.*, 2007), xerophytic (Suryanarayanan *et al.*, 2005) and

aquatic environment (Sraj-Krzic *et al.*, 2006). Medicinal plants are reported to harbour endophytes (Strobel and Daisy, 2003) which in turn provide protection to their host from infection agents and also provide adaptability to survive in adverse environmental condition. It is therefore important to determine the endophytic fungi diversity of medicinal plants. Recently the knowledge about endophyte biodiversity is becoming more apparent.

Centella asiatica (CA) is a very important medicinal herb used in the orient (Bown, 1995), which is also becoming popular in the West (Chevallier, 1996). It is a tasteless, odourless plant that thrives in and around water. It has small fan-shaped green leaves with white or light purple-to-pink or white flowers and it bears small oval fruit. The whole plant is used for medicinal purposes (Singh and Singh, 2002). It is widely used as a blood purifier as well as for treating high blood pressure, for memory enhancement and promoting longevity. In Ayurveda, *Centella asiatica* is one of the main herbs for revitalizing the nerves and brain cells.

Melia azedarach, is a species of deciduous tree in the mahogany family, Meliaceae, that is an evergreen tree, cultivated in various parts of the Indian subcontinents. Neem has a long history of use in the traditional medical systems of

*Corresponding author: Shijila Rani A.S

Department of Botany and Microbiology, AVVM Sri Pushpam College (Autonomous), Poondi, 613 503, Thanjavur

India (Ayurvedic, Unani-Tibb). Leaves have been used as a natural insecticide to keep with stored food, but must not be eaten as they are highly poisonous. A diluted infusion of leaves and trees has been used in the past to induce uterus relaxation (Russell *et al.*, 1997). Extracts from neem leaf, seed, and bark possess a wide spectrum of antibacterial action (Biswas *et al.*, 2002; Mahfuzul *et al.*, 2007). Although the fruits are poisonous part of the tree, they have been used traditionally for the treatment of a variety of diseases, especially dermatitis and rubella (Kim *et al.*, 1999).

MATERIALS AND METHODS

Sample collection

Collection, identification and authentication of leaf materials of *Melia azedarach* L. and *Centella asiatica* L. were collected from Thanjavur District, (the latitude in 10.6590°N, and longitude in 79.2451°E), Tamilnadu, India. The plants were identified by taxonomist for St. Joesph's College, Triuchirappalli (Fig. 1).



Fig 1 Map showing the study site

Isolation of endophytes

The endophytic fungi were isolated from leaves of *Melia azedarach* L. and *Centella asiatica* L. as described by Strobel and Daisy (2003). In first instance, the leaves were cleaned by thorough washing in running tap water, followed by deionized (DI) water. Clean leaves were cut into small pieces were sterilized in series of solution: 70% ethanol; 1% sodium hypochlorite (v/v); again 70% ethanol for 1 min each. Finally they were again rinsed with sterile distilled water two times. After surface sterilization, the tissues were dried on blotting sheets and cut into 1 cm square pieces. These sterile small pieces were placed on Czapek dextrose agar (CDA) plates containing streptomycin ($250 \mu\text{g mL}^{-1}$) to inhibit the bacterial growth.

The plates were incubated at $25 \pm 2^\circ\text{C}$ after wrapping with parafilm and observed daily. The fungal mycelia which started growing from the tissues were sub-cultured on new CDA plates. These endophytes were stored in paraffin at 4°C .

Colony frequency

Colonization frequency (CF %) was calculated as described by Suryanarayanan *et al.* (2003) and Photita (2001). The colonies appearing on petriplates were sub-cultured into the tube containing Czapek dextrose agar medium for identification. Morphological identification was done according to the standard taxonomic key included colony diameter, texture, color and the dimensions and morphology of hyphae and conidia.

$$\text{Percentage of colony} = \frac{\text{No. of colonies in individual segment}}{\text{Total number of colonies in the sample}} \times 100$$

$$\text{Percentage of CF} = \frac{\text{number of segment colonized by fungi}}{\text{total number of segment observed}} \times 100$$

Diversity index

The species diversity of the endophytic fungi was measured using the diversity indices with species richness and relative species abundance. The species evenness, which assesses the contribution of the individuals to the community also calculated

The diversity indices used for the isolates were the Shannon–Wiener index (H) (formula: $[(n_i/n) \ln(n_i/n)]$), Simpson's diversity index (1–D) (formula: $1 - [D = (n_i/n)^2]$) and Simpson's dominance index (D), where n_i is the number of distinct species (i) and (n) is the abundance of each species in the community. The equitability was calculated by the following formula: $E = H/H_{\text{max}}$, where H is the Shannon–Wiener index and H_{max} , is the \ln of n. The diversity analyses were performed the MS Excel program (Hammer *et al.*, 2001).

RESULTS

In the present study, eleven isolates were obtained from seven tissue fragments from two medicinal plant species. 10 different types of genera were isolated, the maximum number of colonies of *Absidia repens* and *Aspergillus terreus* were observed in *M.azedarach* and *A.terreus* in *C.asiatica* leaves fragment. The number of colony, percentage of colony and percentage of colony frequency were recorded (Table 1).

Table 1 Isolation of endophytic fungi from the leaf samples

S.No	Name of the fungi	<i>M. azedarach</i>			<i>C.asiatica</i>		
		No. of colony	% of CP	Colony Frequency	No. of colony	% of CP	Colony frequency
1	<i>Absidia repens</i>	5	20	71.42	1	6.25	20
2	<i>Alternaria</i> sp.	1	4	14.28	1	6.25	20
3	<i>Aspergillus flavus</i>	4	16	57.14	2	12.5	40
4	<i>A.terreus</i>	5	20	71.42	4	25	80
5	<i>Chaetomium globosum</i>	1	4	14.28	1	6.25	20
6	<i>Curvularia geniculata</i>	1	4	14.28	-	0	0
7	<i>C.lunata</i>	-	0	0	1	6.25	20
8	<i>Curvularia</i> sp.	2	8	28.57	2	12.5	40
9	<i>Fusarium solani</i>	3	12	42.85	3	18.75	60
10	<i>Humicola</i> sp.	2	8	28.57	1	6.25	20
11	<i>Melanospora</i> sp.	1	4	14.28	-	0	0
12	Total	25	100	357.14	16	100	320

Endophytic fungal diversity is measured in terms of Shannon, Margalef's index, Menhinick's species richness, Margalef's Diversity, McIntosh's Evenness, Simpson and Pielou's evenness and Fisher's alpha indices. The diversity indices were calculated in the endophytic fungi of both the plants. The highest species richness present in *M.azedarach* when compared to *C.asiatica*. The Margalef's Diversity is 14.427 in *C.asiatica* leaves and 12.427 in *M.azedarach* (Table 2).

Table 2 Diversity indices of isolated endophytic fungi

S.No	Diversity indices	<i>M.azedarach</i>	<i>C.asiatica</i>
1	No. of Species	25	16
2	No. of individuals	10	9
3	Genera	10	8
4	Margalef's index (d)	6.463	4.039
5	Menhinick's species richness (SR)	5.735	3.671
6	Pielou's Evenness (J)	0.086	0.081
7	McIntosh's Evenness (McE)	0.141	0.188
8	Shannon's Diversity (H)	0.277	0.224
9	Margalef's Diversity (Dmg)	12.427	14.427
10	Fisher's alpha	6.17	8.5

DISCUSSION

Endophytic microorganism especially fungi established a symbiotic relationship with plants while living inside the plant and synthesize metabolites that often helps their host to survive and adapt in adverse environment (Clay and Schardl, 2002). In addition, they are the group of organism with very good potential for application in plant improvement and disease control (Sandhu *et al.*, 2014). A study of endophytic biodiversity of the temporal variation in *Plumeria rubra* leaves was conducted by the Suryanarayanan and Thennarasan (2004). In the similar study 2 different genera of *Aspergillus* (6.6%) and *Nigrospora* (3.3%) the colonization frequency were isolate from *Melia azedarach*. This is slightly less than the above cited report by Suryanarayanan and Thennarasan (2004). During the present study, *Absidia* sp., *Alternaria* sp., *Aspergillus* sp., *Chaetomium* sp., *Curvularia* sp., *Fusarium* sp., *Humicola* sp. and *Melanospora* sp. were isolated from *Melia azedarach* and *Centella asiatica*. All the species belong to Hyphomycetes class. These class fungi were largely prevalent and showed the resistance too many pathogens. The genus *Aspergillus* found most common in plant.

Similar to the findings reported previously in *Centella asiatica* leaves, the higher colonization of endophytic fungi in mature leaf as compared to young leaf over all the three seasons suggests that mature leaf provides more favourable environment for the establishment of endophytic fungi (Gupta and Preeti, 2017). Nascimento *et al.*, (2015) reported *Phaeoramularia calotropidis* (63.5%) as dominant endophyte in *Calotropis procera* leaves.

The diversity of the endophytic fungal populations isolated from leaves of *C.asiatica* and *Melia azedarach* was evaluated by various indices such as Fisher alpha diversity index (α), Shannon index H, Simpson's diversity index (1-D) and Margalef richness index (Dmg) (Table 2). Similarly, Menhinick's index is similar, conceptually, to Margalef's richness index for analyzing species richness. The assume diversity values from each of the two indices showed similar patterns in these results, because of the endophytes, Shannon's diversity index (H') (Pielou's, 1975) and Simpson's diversity index (D) (Simpson, 1949) were not used. The species richness

(Dmn) was highest in the leaves (4.24) followed by the flowers (2.82) and roots (1.41) (Katoch *et al.*, 2017).

Acknowledgement

The authors thank the principal of A.V.V.M Sri Pushpam College (Autonomous), Poondi, Thanjavur and the Director, Indian Biotrack Research Institute, Thanjavur for extending the Lab facilities to carry out the work.

Reference

- Arnold, AE. and Herre, EA., 2003. Canopy cover and leaf age affect colonization by tropical fungal endophytes: Ecological pattern and process in *Theobroma cacao* (Malvaceae). *Mycologia*; 95 (3):388-398.
- Biswas, K., Chattopadhyay, T., Banerjee, R.K. and Bandyopadhyay, U., 2002. Biological activities and medicinal properties of neem (*Azadirachta indica*). *Curr. Sci.*; 82: 1336-1345.
- Bown, D., 1995. Encyclopaedia of Herbs and their Uses. London: Dorling Kindersley: 361-365.
- Chevallier, A., 1996. The Encyclopedia of Medicinal Plants. London: Dorling Kindersley: 257.
- Clay, K. and Schardl, C., 2002. Evolution origins and ecological consequences of endophytes symbiosis with grasses. *American Naturalist.*; 160: 99-127.
- Ganley, RJ., Brunfeld, SJ. and Newcombe, G., 2007. A community of unknown, endophytic fungi in western white pine. *Proc Nat Acad Sci USA*: 101:10107-10112.
- Gupta, S. and Preeti C., 2017. Foliar Endophytic Diversity of *Centella asiatica* (L.) Urban in Relation to Different Seasons and Leaf Age. *Int.J.Curr.Microbiol.App.Sci.*; 6(6): 468-477.
- Hammer, O., Harper, DAT. and Ryan, PD., 2001. PAST: Paleontological statistics software package for education and data analysis. *Paleontologica Electronica.*; 4(9):178.
- Katoch, M., Phull, S., Vaid, S. and Singh, S., 2017. Diversity, Phylogeny, anticancer and antimicrobial potential of fungal endophytes associated with *Monarda citriodora* L. *BMC Microbiology*; 17:44.
- Kim, M., Kim, SK., Park, BN., Lee, KH., Min, GH., Seoh, J.Y., Park CG., Hwang ES., Cha C.Y. and Kook, YH., 1999. Antiviral effects of 28-deacetylSENDANIN on herpes simplex virus-1 replication. *Phytochemistry*; 43:103-112.
- Mahfuzul, H.M.D., Bari, M.L., Inatsu, Y., Vijay, K.J. and Kawamoto, S., 2007. Antibacterial Activity of Guava (*Psidium guajava* L.) and Neem (*Azadirachta indica* A. Juss.) Extracts Against Foodborne Pathogens and Spoilage Bacteria. *Foodborne pathogens disease.*; 4(4): 481-488.
- Mohali, S., Burgess, TI. and Wingfield, MN., 2005. Diversity and host association of the tropical tree endophytic *Lasiodiplodia theobromae* revealed using simple sequence repeat markers. *Forest pathology.*; 35 (6):385-396.
- Nascimento, T.L., Oki, Y., Lima, D.M.M., Almeida-Cortez, J.S., Wilson Fernandes, G. and Souza-Motta, C.M., 2015. Biodiversity of endophytic fungi in different leaf ages of *Calotropis procera* and their antimicrobial activity. *Fungal Ecol.*; 14: 79-86.

- Neto, P.A.S.P., Azevedo, J.L. and Araujo, W.L., 2002. Microorganisms endofíticos: interacao complaints e potencial biotecnológico. *Biotecnol Ciencia Desenvolvimento*; 29:62–76.
- Photita, W, Lumyong, S, Lumyong, P. and Hyde, K.D., 2001. Endophytic fungi of wild banana (*Musa acuminata*) at doi Suthep Pui National Park, Thailand. *Mycol Res.*, 105: 1508-1513.
- Pielou, E.C., 1975. Ecological diversity. New York: John Wiley & Sons.
- Prompttha, I., Lumyong, S., Dhanasekaran, V., McKenzie, E.H., Hyde, K.D. and Jeewon, R.A., 2007. Phylogenetic evaluation of whether endophytes become saprotrophs at host senescence. *Microb Ecol*; 53(4):579–90.
- Redman, R.S., Sheehan, K.B., Stout, R.G., Rodriguez, R.J. And Henson, J.M., 2002. Thermotolerance generated by plant/fungal symbiosis. *Science*; 298:1581
- Russell, A.B., Hardin, J.W. and Grand, L., 1997. *Melia azedarach*. In: Poisonous Plants of North Carolina.
- Sandhu, S.S., Kumar, S. and Aharwal, R.P., 2014. Isolation and Identification of Endophytic Fungi from *Ricinus communis* Linn. and their antibacterial activity. *Inter. J. Rese. In phar. And chemi.*, 4(3): 611-618.
- Sieber, T.N., 2007. Endophytic fungi in forest trees: are they mutualists. *Fungal Biol Rev*; 21(2-3):75-89.
- Simpson, E.H., 1949. Measurement of diversity. *Nature*;163: 688.
- Singh, P. and Singh, J.S., 2002. Recruitment and competitive interaction between ramets and seedlings in a perennial medicinal herb, *Centella asiatica*. *Basic Appl Ecol*; 3:65–76.
- Sraj-Krzic, N., Pongrac, P., Klemenc, M., Kladnik, A., Regvar, M. and Gagerscik, A., 2006. Mycorrhizal colonization in plants from intermittent aquatic habitats. *Aquatic Botany.*; 85 (4): 340-346.
- Stone, J.K., Polishook, J.D. and White, J.F., 2004. Endophytic fungi. In: Biodiversity of fungi. Inventory and monitoring methods (Mueller G.M., Bills G.F., Foster M.S., eds). Elsevier Academic Press, San Diego, USA: 241-270.
- Strobel, G.A. and Daisy, B., 2003. Bioprospecting for microbial endophytes and their natural products. *Microbiol Mol Biol Rev.*; 67: 491-502.
- Strobel, G.A., 2000. Microbial gifts from rain forests. *Canadian Journal of plant pathology*; 24:14- 20.
- Suryanarayanan, T.S. and Thennarasan, S., 2004. Temporal variation in endophyte assemblages of *Plumeria rubra* leaves. *Fungal Diversity.*; 15: 197-204.
- Suryanarayanan, T.S., Venkatesan, G. and Murali, T.S., 2003. Endophytic fungal communities in leaves of tropical forest trees: Diversity and distribution patterns. *Current Science*, 85(4):489 - 492.
- Suryanarayanan, T.S., Wittlinger, S.K.J. and Faeth, S.H., 2005. Endophytic fungi associated with cacti in Arizona. *Mycological Research*; 109 (5):635-639.

How to cite this article:

Shijila Rani A.S., Ambikapathy, V and Panneerselvam A.2018, Diversity of endophytic fungi from leaves of *Centella asiatica* L. and *Melia azedarach* L. *Int J Recent Sci Res.* 9(1), pp. 23083-23086. DOI: <http://dx.doi.org/10.24327/ijrsr.2018.0901.1394>
