



ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

International Journal of Recent Scientific Research
Vol. 7, Issue, 10, pp. 13955-13958, October, 2016

**International Journal of
Recent Scientific
Research**

Research Article

STUDY OF THROMBOCYTOPAENIA IN MALARIA IN RURAL POPULATION OF RAYALASEEMA, A.P

**Abdul Razaak., Ilyas Ahmed., Feroz Khan*., Raghavaram N., Anil Kumar.,
Srujana and PrabhakarRao R**

Santhiram Medical College, Nandyal, (Kurnool Dt) Andhra Pradesh. Pin – 518 501

ARTICLE INFO

Article History:

Received 18th July, 2016
Received in revised form 10th August, 2016
Accepted 06th September, 2016
Published online 28th October, 2016

Key Words:

Thrombocytopenia, Falciparum malaria,
Vivax malaria

ABSTRACT

Malaria is a major health problem in India. It can be associated with many complications. Thrombocytopenia is a common feature in malaria. In Patients having fever with thrombocytopenia, malaria should be on top of differential diagnosis. This was a study of admitted patients in Nandyal, rural area of Rayalaseema region of Andhra Pradesh, diagnosed with malaria and they were assessed for thrombocytopenia using automated quantitative D3 Analyzer. The data was analyzed using SPSS version 16.0. The results were obtained in numbers and percentages. Data was expressed as means with standard deviation. Viral hemorrhagic fevers can also present as fever but in these cases the complete blood counts has other features such as high hematocrit and leucopenia along with thrombocytopenia. Our finding can have therapeutic implications in the context of avoiding unnecessary platelet transfusion with the relatively more benign course in vivax malaria. In Our study the commonest cause of thrombocytopenia in malaria was vivax malaria.

Copyright © Abdul Razaak et al., 2016, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Malaria is commonly associated with various degrees of hematological complications like anemia and thrombocytopenia. The anemia is usually due to varied reasons ranging from hemolysis to comorbidities like parasitic infections, folate, iron, and vitamin B12 deficiencies in endemic areas, antimalarials and further complicated by the coexistence of thalassemia and other haemoglobinopathies¹. Anemia and thrombocytopenia are the most frequent malaria-associated hematological complications². The hematopoietic response is, however, also somewhat blunted, and there is reduced platelet count and sometimes reduced WBC counts as well. Thrombocytopenia is a very common association of malaria. In the past it was thought that malaria is rarely associated with clinical features of thrombocytopenia like bleeding disorders and is usually an incidental finding on blood testing. Recent scientific evidences have invalidated this anecdote. It is pertinent that the finding of thrombocytopenia in patient may be an indication for a thorough lookout into the blood smear to rule out malaria as the cause. This fact is especially important in the workup for thrombocytopenia in febrile patients. Thrombocytopenia may be associated with bleeding tendency which is one of the important severe manifestations of P. Falciparum malaria. The presence of thrombocytopenia in acute febrile travelers returning from tropical areas has become a highly sensitive clinical marker for

malaria diagnosis³. A number of observational studies have confirmed the association of thrombocytopenia to malaria but till date the cause of thrombocytopenia is poorly understood. The speculated mechanisms leading to thrombocytopenia are coagulation disturbances, splenomegaly, bone marrow alterations, antibody-mediated platelet destruction, oxidative stress, and the role of platelets as cofactors in triggering severe malaria⁴⁻⁶. In an animal model of cerebral malaria, platelet's role has been implicated. In autopsy studies of children dying of cerebral malaria, clumps of platelets with or without infected RBC's have been found in the brain capillaries^{7,8}. The platelet counts usually are normalized quickly in a matter of few weeks when malaria is treated successfully.

MATERIAL AND METHODS

It was a prospective observational study conducted in all patients of fever with thrombocytopenia presenting to the Medical Unit of Santhiram Medical College and General Hospital during November 2013 to November 2015 were included in the study.

The blood counts were visually confirmed by a clinical hematologist after the counts were done by an auto analyzer machine. The thick and thin slides were prepared according to the WHO guidelines and studied by a hematologist. Thick smears using Geimsa stain and thin smears using Wright stain were used and the patient was labeled as nonmalaria only if

*Corresponding author: **Feroz Khan**

Santhiram Medical College, Nandyal, (Kurnool Dt) Andhra Pradesh. Pin – 518 501

three consecutive smears were negative. Thus these patients were divided further into two groups: malaria and non-malaria group. The non-malaria group was further evaluated for other causes of thrombocytopenia which was not part of this study. Laboratory tests done on these patients consisted of full blood counts, liver enzymes, renal function, vitamin B12 levels, folic acid levels, erythrocyte sedimentation rate, and peripheral blood smear. In undiagnosed cases and where suspicion of immune thrombocytopenia was high, bone marrow biopsy was done to differentiate whether the low platelet count is due to decreased production or peripheral destruction.

Statistical calculation was done using Word Excel and SPSS version 11 for diagnostic accuracy between the two groups, that is, between platelet count from 25,000/dL to 50,000/dL, 50,000/dL to 75,000/dL and 75,000/dL to 150,000/dL using the Wilson score with 95% confidence interval.

Observations

Of the total of 556 patients with fever and thrombocytopenia, 199 patients (53%) proved to be suffering from malaria. Of them 199 patients (68%) had falciparum malaria while 95 patients (32%) had vivax infection. Of these 121 patients, platelet counts ranged between 25,000 and 150,000/dL with a mean value of 101,000/dL (S D ± 4 7, 5 0 0) and a median of 75,000/dL. Of the 107 patients who were not suffering from malaria, the counts ranged between 10,000 and 150,000/dL with a mean value of 58,000/dL (S D ± 5 4, 0 0 0) and median of 50,000/dL..

The results are shown in table 1 and table 2

Table – 1

Malaria Cases Types	
Falciparum	199
Vivax	95





Table-2

Platelet Counts In Malaria Pts And Not Malaria Pts	
With Malaria	25,000 TO 1.5 L
Without Malaria	10,000 TO 1.5 L




Platelet counts of 75,000 to 150,000/dL are defined as grade 1 thrombocytopenia, 50,000 to <75,000/dL as grade 2, 25,000 to <50,000/dL as grade 3, and below 25,000/dL as grade 4 thrombocytopenia.

The results are shown in table 3

Table 3


Grading Of Thrombocytopenia As Per Platelet Count	
Grade 1	75,000 TO 1.5 L
Grade 2	50,000 TO 0.75 L
Grade 3	25,000 TO 0.50 L
Grade 4	<25,000



Thrombocytopenia was thus divided into these groups. The numbers of patients who had and had not malaria after the evaluation are mentioned in Table 4.

Table 4

Fever With Thrombocytopenia	
With Malaria	294
Without Malaria	262



The accuracy parameters between different levels of platelet counts are given in Table 5

Wilson Score

Level 1	50,000 TO 75,000
Level 2	75,000 TO 1.5 L




Table 2: Accuracy parameters between level 1 (platelets from 25,000 to 50,000 dL) and level 2 (platelets from 50,000 to 75,000 dL) using Wilson score.

DISCUSSION

Thrombocytopenia is associated with malaria and is usually mild to moderate. It may however be symptomatic and severe⁹⁻¹⁰. Fifty-three percent of patients with malaria showing thrombocytopenia in our study is close to others reporting low platelets as 57% [9] and 48%¹⁰. In Liberia, Mahmood¹¹ and Yasir studied a total of 145 patients who had *P. falciparum* malaria. Out of these 109 (75.18%) had thrombocytopenia. The sensitivity of the platelet count was considered as a predictor of malaria was 80.11% while specificity was 81.36%. The positive predictive value was 63.87% and the negative predictive value was 90.86%. Another study has reported 60% sensitivity and 88% specificity of thrombocytopenia for malaria diagnosis in acute febrile patients¹². In another important study from India, Patel *et al.* reports the sensitivity of thrombocytopenia together with the acute febrile syndrome as 100% for malaria diagnosis, with a specificity of 70%, a positive predictive value of 86% and a negative predictive value of 100%¹³. In our study we have calculated the sensitivity, specificity, positive and negative predictive value and diagnostic accuracy at different levels of platelet counts and have concluded that the sensitivity and negative predictive value is high at low platelet counts (78% sensitivity and 67% negative predictive value 40% and 58% resp.) while the specificity and positive predictive value is high at relatively higher platelet counts (94% specificity and 89% positive predictive value against 50% and 64% resp.). The diagnostic accuracy remains almost the same at both counts (around 65%). Mahmood and Yasir¹¹ concluded an extended search for malarial parasite in patients having thrombocytopenia on smear. Mild-to-severe thrombocytopenia observed in hospitalized patients was considered enough to alert the possibility of malarial infection, as *P. falciparum* was found to be common species in these patients. Our study also connotes that *falciparum* malaria is more common at lower platelet counts as compared to *vivax* infection and overall the chances of finding *falciparum* malaria are almost twice than that of finding *vivax* malaria in thrombocytopenic patients. It is a general consensus that thrombocytopenia is very common in malaria^{14,15} and previously it was believed that it is more common in *F. malaria*. Recent studies have shown that thrombocytopenia is equally or even more common in *P. vivax* malaria contrary to the popular belief that it may be observed in *P. falciparum* malaria¹⁶⁻²³. More recent data in India has shown how thrombocytopenia exhibited a heightened frequency and severity among patients with *P. vivax* infection²⁴. Recent studies conducted from the Indian subcontinent have found significant thrombocytopenia in *P. vivax* malaria^{25,26}. Similar results have been reported from Qatar and Venezuela^{27,28}. Studies from Brazil have shown a similar trend²⁹. A recent study from Iran confirms that they are getting more cases of thrombocytopenia due to *P. vivax* than *Falciparum* and attributes this to the possible development of a new genotype of *P. vivax*³⁰. Though we have found significant thrombocytopenia in more than half of our patients with malaria, the fact that it was seemingly more common in *P. falciparum* malaria needs to be validated by larger studies.

CONCLUSIONS

The presence of thrombocytopenia may be a predictor of malaria in adult population. Malaria should be a consideration in all patients with low platelets and after excluding this common and easily treatable cause, further evaluation of thrombocytopenia should be undertaken.

References

1. K. Ghosh and K. Ghosh, "Pathogenesis of anemia in malaria: a concise review," *Parasitology Research*, vol. 101, no. 6, pp. 1463–1469, 2007. View at Publisher · View at Google Scholar · View at Scopus
2. S. N. Wickramasinghe and S. H. Abdalla, "Blood and bone marrow changes in malaria," *Bailliere's Best Practice and Research in Clinical Haematology*, vol. 13, no. 2, pp. 277–299, 2000. View at Publisher · View at Google Scholar · View at Scopus
3. V. D'Acromont, P. Landry, I. Mueller, A. Pécoud, and B. Genton, "Clinical and laboratory predictors of imported malaria in an outpatient setting: an aid to medical decision making in returning travelers with fever," *American Journal of Tropical Medicine and Hygiene*, vol. 66, no. 5, pp. 481–486, 2002. View at Google Scholar · View at Scopus
4. S. S. Jamal Khan, F. R. Khan, M. Usman, and S. Zahid, "Malaria can lead to thrombocytopenia," *Rawal Medical Journal*, vol. 33, no. 2, pp. 183–185, 2008. View at Google Scholar · View at Scopus
5. T. S. Faseela, R. A. Roche, K. B. Anita, C. S. Malli, and Y. Rai, "Diagnostic value of platelet count in malaria," *Journal of Clinical and Diagnostic Research*, vol. 5, no. 3, pp. 464–466, 2011. View at Google Scholar · View at Scopus
6. A. Rasheed, S. Saeed, and S. A. Khan, "Platelet count in malaria," *Pakistan Journal of Phytopathology*, vol. 19, no. 3, pp. 86–88, 2008. View at Google Scholar
7. G. E. Grau, C. D. Mackenzie, R. A. Carr *et al.*, "Platelet accumulation in brain microvessels in fatal pediatric cerebral malaria," *Journal of Infectious Diseases*, vol. 187, no. 3, pp. 461–466, 2003. View at Publisher · View at Google Scholar · View at Scopus
8. A. Pain, D. J. P. Ferguson, O. Kai *et al.*, "Platelet-mediated clumping of *Plasmodium falciparum* erythrocytes is a common adhesive phenotype and is associated with severe malaria," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 98, no. 4, pp. 1805–1810, 2001. View at Publisher · View at Google Scholar · View at Scopus
9. G. A. Niazi, "Haematological aspect of malaria in a population based hospital, Saudi Arabia," *Journal of the Egyptian Society of Parasitology*, vol. 25, no. 3, pp. 787–793, 1995. View at Google Scholar · View at Scopus
10. A. Mert, R. Ozaras, F. Tabak, M. Bilir, R. Ozturk, and Y. Aktuglu, "Malaria in Turkey: a review of 33 cases," *European Journal of Epidemiology*, vol. 18, no. 6, pp. 579–582, 2003. View at Publisher · View at Google Scholar · View at Scopus
11. A. Mahmood and M. Yasir, "Thrombocytopenia: a predictor of malaria among febrile patients in Liberia,"

- Infectious Diseases Journal*, vol. 14, pp. 41–44, 2008. View at Google Scholar
12. T. B. Lathia and R. Joshi, “Can hematological parameters discriminate malaria from nonmalarious acute febrile illness in the tropics?” *Indian Journal of Medical Sciences*, vol. 58, no. 6, pp. 239–244, 2004. View at Google Scholar · View at Scopus
 13. U. Patel, G. Gandhi, S. Friedman, and S. Niranjana, “Thrombocytopenia in malaria,” *Journal of the National Medical Association*, vol. 96, no. 9, pp. 1212–1214, 2004. View at Google Scholar · View at Scopus
 14. M. N. Akhtar, S. Jamil, S. I. Amjad, A. R. Butt, and M. Farooq, “Association of malaria with thrombocytopenia,” *Annals of King Edward Medical College*, vol. 11, pp. 536–537, 2005. View at Google Scholar
 15. Z. U. Rehman, M. Alam, A. Mahmood, A. Mubarik, A. Sattar, and K. A. Karamat, “Thrombocytopenia in acute malarial infection,” *Pakistan Journal of Pathology*, vol. 10, pp. 9–11, 1999. View at Google Scholar
 16. A. Aggarwal, S. Rath, and Shashiraj, “Plasmodium vivax malaria presenting with severe thrombocytopenia,” *Journal of Tropical Pediatrics*, vol. 51, no. 2, pp. 120–121, 2005. View at Publisher · View at Google Scholar · View at Scopus
 17. N. M. Anstey, B. J. Currie, and M. E. Dyer, “Profound thrombocytopenia due to Plasmodium vivax malaria,” *Australian and New Zealand Journal of Medicine*, vol. 22, no. 2, pp. 169–170, 1992. View at Google Scholar · View at Scopus
 18. V. Bhatia and J. Bhatia, “Severe thrombocytopenia with bleeding manifestations in two children secondary to Plasmodium vivax,” *Platelets*, vol. 21, no. 4, pp. 307–309, 2010. View at Publisher · View at Google Scholar · View at Scopus
 19. R. Harish and S. Gupta, “Plasmodium vivax malaria presenting with severe thrombocytopenia, cerebral complications and hydrocephalus,” *Indian Journal of Pediatrics*, vol. 76, no. 5, pp. 551–552, 2009. View at Publisher · View at Google Scholar · View at Scopus
 20. U. M. Jadhav, V. S. Patkar, and N. N. Kadam, “Thrombocytopenia in malaria—correlation with type and severity of malaria,” *Journal of Association of Physicians of India*, vol. 52, pp. 615–618, 2004. View at Google Scholar · View at Scopus
 21. A. Kakar, S. Bhoi, V. Prakash, and S. Kakar, “Profound thrombocytopenia in Plasmodium vivax malaria,” *Diagnostic Microbiology and Infectious Disease*, vol. 35, no. 3, pp. 243–244, 1999. View at Publisher · View at Google Scholar · View at Scopus
 22. D. Kaur, V. Wasir, S. Gulati, and A. Bagga, “Unusual presentation of Plasmodium vivax malaria with severe thrombocytopenia and acute renal failure,” *Journal of Tropical Pediatrics*, vol. 53, no. 3, pp. 210–212, 2007. View at Publisher · View at Google Scholar · View at Scopus
 23. A. Kumar and Shashirekha, “Thrombocytopenia—an indicator of acute vivax malaria,” *Indian Journal of Pathology and Microbiology*, vol. 49, no. 4, pp. 505–508, 2006. View at Google Scholar · View at Scopus
 24. D. K. Kochar, A. Das, A. Kochar et al., “Thrombocytopenia in Plasmodium falciparum, Plasmodium vivax and mixed infection malaria: a study from Bikaner (Northwestern India),” *Platelets*, vol. 21, no. 8, pp. 623–627, 2010. View at Publisher · View at Google Scholar · View at Scopus
 25. S. Srivastava, S. Ahmad, N. Shirazi, S. Kumar Verma, and P. Puri, “Retrospective analysis of vivax malaria patients presenting to tertiary referral centre of Uttarakhand,” *Acta Tropica*, vol. 117, no. 2, pp. 82–85, 2011. View at Publisher · View at Google Scholar · View at Scopus
 26. P. George and L. M. Alexander, “A study on the clinical profile of complicated Plasmodium vivax mono-infections,” *Asian Pacific Journal of Tropical Medicine*, vol. 3, no. 7, pp. 560–562, 2010. View at Publisher · View at Google Scholar · View at Scopus
 27. F. Yousef Khan, A. K. Lutof, M. A. Yassin et al., “Imported malaria in Qatar: a one year hospital-based study in 2005,” *Travel Medicine and Infectious Disease*, vol. 7, no. 2, pp. 111–117, 2009. View at Publisher · View at Google Scholar · View at Scopus
 28. B. González, H. Rodulfo, M. De Donato, M. Berrizbeitia, C. Gómez, and L. González, “Hematologic variations in patient with malaria caused by Plasmodium vivax before, during and after treatment,” *Investigacion Clinica*, vol. 50, no. 2, pp. 187–201, 2009. View at Google Scholar · View at Scopus
 29. S. B. R. Silva, Avaliação da frequência e dos fatores associados à plaquetopenia causada pelo Plasmodium vivax [MSc Thesis], Universidade Federal do Mato Grosso, Mato Grosso, Brazil, 2009.
 30. Metanat and B. Sharifi-Mood, “Malaria vivax and severe thrombocytopenia in Iran,” *Iranian Journal of Parasitology*, vol. 5, no. 3, pp. 69–70, 2010. View at Google Scholar · View at Scopus

How to cite this article:

Abdul Razaak et al., Study of Thrombocytopaenia in Malaria In Rural Population of Rayalaseema, A.P. *Int J Recent Sci Res.* 7(10), pp. 13955-13958.