



## CHRONIC SUPPURATIVE OTITIS MEDIA INFECTIONS: BACTERIAL PROFILE AND DEMOGRAPHIC FEATURES AMONG PATIENTS ATTENDING TERTIARY CARE HOSPITAL

Jyoti R. Hundekar\*<sup>1</sup>, Surekha, Y.A.<sup>2</sup>, S. Sadiqa Begum<sup>3</sup> and Dr Sharada C Metgud

<sup>1,4</sup>Department of Microbiology, KLE Jagadguru Gangadhar Mahaswamigalu Moorsavirmath Medical College and Hospital, Hubballi- 580028, Karnataka, India

<sup>2</sup>Department of Microbiology, VIMS, Ballari.

<sup>3</sup>Department of Microbiology, Subbaiah Institute of Medical Science, Shivamogga, Karnataka, India

DOI: <http://dx.doi.org/10.24327/ijrsr.20251601.009>

### ARTICLE INFO

#### Article History:

Received 16<sup>th</sup> December, 2024

Received in revised form 31<sup>st</sup> December 2024

Accepted 22<sup>nd</sup> January 2025

Published online 28<sup>th</sup> January, 2025

#### Key words:

Chronic Suppurative Otitis Media Infections,  
Bacterial Isolates, Demographic Features

### ABSTRACT

**Introduction:** Chronic suppurative otitis media (CSOM) can cause intracranial and extracranial complications. Reducing its prevalence may have both immediate and long-term health advantages. Understanding the local microbial flora in CSOM is critical before beginning empirical therapy. Our study sought to evaluate the link between bacterial profiles and demographic features in patients admitted to tertiary care institutions with persistent suppurative otitis media infections. **Materials & Methods:** The study was conducted in the Department of Microbiology at the Vijayanagar Institute of Medical Sciences in Ballari. Random selection was used to choose 109 CSOM cases who were admitted to ENT wards and visited outpatient departments. A proforma was used to record each patient's age, sex, address, complete clinical information, and risk factors. Utilizing conventional microbiological techniques, bacterial isolates were obtained. **Result:** The largest percentages of patients (28.4%) were in the 1 to 11 years age range. Compared to women (36%), men were more impacted (64%). The prevalence of CSOM patients was higher in rural areas (66%) than in urban areas (34%). 86 of the 129 ear samples that tested positive for culture contained 100 bacterial isolates; *Staphylococcus aureus* 36 (36%) was the most frequently isolated pathogen, followed by *Pseudomonas spp.* **Conclusion,** CSOM affects both adults and children. CSOM is commonly dismissed as a minor public health concern. The incidence and morbidity of this disorder can be reduced with the support of health awareness initiatives, improved health education, and easy access to medical facilities.

Copyright© The author(s) 2025, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

### INTRODUCTION

Chronic Suppurative Otitis Media (CSOM) is an infection of the middle ear that lasts more than 3 months and is accompanied by tympanic membrane perforation<sup>1</sup>. The annual global incidence is estimated at 4.76 per 1000 people per year, totalling 31 million cases, with 22.6% of these cases arising in children under 5 years old<sup>2</sup>. The highest incidence is found in the Western Pacific and South East Asian regions<sup>3</sup>. Infection can spread from the middle ear to vital structures

leading to complications. The reported overall extracranial and intracranial complication rate in CSOM varies from 0.7% to 3.2%. It was found to be the single major cause for conductive deafness (66.3%) and is also responsible for 1.5% of speech disorders. Hearing loss and ear diseases in India according to WHO, CSOM accounts about 5.2%.

CSOM has garnered significant focus, not only due to its frequent occurrence and long-lasting nature but also because of concerns like bacterial resistance and ototoxicity associated with both topical and systemic antibiotics. The wide spread use of antibiotics has precipitated the emergence of multiple resistant strains of bacteria which can produce both primary and post-operative ear infections. The WHO highlights the need to clearly establish risk factors for the development of CSOM<sup>3</sup>. Determining the risk factors for the development of

\*Corresponding author: Jyoti R. Hundekar

Assistant professor, Department of Microbiology, KLE Jagadguru Gangadhar Mahaswamigalu Moorsavirmath Medical College and Hospital, Hubballi- 580028, Karnataka, India

CSOM will aid in the prevention of the disease and lower the related morbidity and mortality.

**MATERIALS AND METHODS**

The present study “Chronic Suppurative Otitis Media infections in relation with Bacteriological profile and Their Demographic Features Among Patients Attending Tertiary Care Hospital” was conducted in the tertiary care hospital, Vijayanagar institute of medical sciences, Ballari, from December 2015 to January 2017. 109 cases with CSOM of all age groups and both sexes attending outpatient department and those admitted in ENT wards were selected randomly for the study.

**Inclusion criteria:** All clinically diagnosed cases of CSOM of all age groups and both sexes attending ENT outpatient department were included.

**History taking and examination:** A proforma was filled for each patient documenting age, sex, address, detailed clinical information, and predisposing factors.


**Collection of sample:** In clinically diagnosed cases of CSOM attending ENT outpatient department were selected randomly. Ears were inspected, then cleaned with spirit to remove contaminant<sup>15</sup>. Ear discharges collected by two sterile swabs under all aseptic precautions<sup>6</sup>. Swabs were sent to microbiology laboratory for further processing<sup>15</sup>. One swab for Gram staining and the other one for an aerobic culture<sup>6</sup>, plating on Blood agar, Macconkey’s agar. All plates were incubated aerobically at 37°C and evaluated at 24 hours to 48 hours. The specific identification of bacterial pathogens was done based on microscopic morphology, staining characteristics, cultural and biochemical properties using standard laboratory procedures.

**RESULTS**

Clinically diagnosed 109 cases of chronic suppurative otitis media attending ENT outpatient department were studied in tertiary care hospital, Vijaynagar institute of Medical sciences, Ballari. Observations made from the study are as follows.

Age (1 year to 66years)	Sex: Male/Female	
	Male (%)	Female (%)
1 to 10 yrs	28.6(%)	11(28.2%)
11 to 20 yrs	27.1(%)	8(20.5%)
21 to 30 yrs	20(%)	7(17.9%)
31 to 40 yrs	7(10%)	3(7.7%)
>41 yrs	10(14.3%)	10(25.6%)

The study group’s subjects range in age from one year to sixty seven years. There were 39 female cases and 70 male cases out of 109 totals. The age range of 1–10 years had the highest incidence of CSOM (28.4%).




**Graph 1** Allocation of patients based on area

38 (35%) of the 109 cases were from cities, while 71 (65%) came from rural areas. The incidence of CSOM was higher in rural than in urban areas.

Ear involved	Frequency	Percentage (%)
Bilateral	20	18
Right	35	32
Left	54	50
Total	109	100

were 35 cases (32%) with a right ear infection, 54 cases (50%) with a left ear infection, and 20 cases (18%) with a bilateral infection.



**Graph 2** Culture results

According to the above table, 20 of the 129 samples were deemed contaminated and were not included in the study. Of the 109 samples, 23 (21%) were culture negative and 86 (79%) were culture positive.

Pattern of Organisms	No of isolates
<b>Gram positive cocci</b>	
<i>Staphylococcus aureus</i>	36(36%)

<i>Coagulase negative staphylococci(CONS)</i>	9 (9%)
<i>Streptococcus spp</i>	4 (4%)
<b>Gram negative organisms</b>	
<i>Pseudomonas aeruginosa</i>	34(34%)
<i>E.coli</i>	4(4%)
<i>Citrobacter freundii</i>	4(4%)
<i>Acinetobacter spp.</i>	4(4%)
<i>Klebsiella spp</i>	2(2%)
<i>Proteus Mirabilis</i>	2(2%)

One hundred bacterial isolates were isolated from 86 of the 129 ear samples that demonstrated culture positive. According to the above table, *Staphylococcus aureus* 36 (36%) was the most often isolated bacterium, followed by *Pseudomonas* spp. 34 (34%), and Coagulase-negative *Staphylococcus* 9 (9%). *Acinetobacter* spp., *E. coli*, and *Citrobacter* spp. are the remaining four (4%). There were two (2% each) *Klebsiella* and *Proteus mirabilis* and one (1% each) *Enterobacter*.

Sex	Culture	
	Positive	Negative
	Number(%)	Number(%)
Female	27(31%)	12(52%)
Male	59(69%)	11(48%)
Total	86(100%)	23(100%)

Of the 70 (64%) samples taken from male patients, 11 (48%) were culture negative and 59 (69%) were culture positive. Twelve (52%) and 27 (31%) of the 39 (36%) samples that were taken from the female patients were culture negative. Male patients in this study exhibited higher levels of cultural positive than female patients.

Age group	Culture result		Total Number (%)
	Positive culture Number (%)	Negative culture Number (%)	
1 to 10 years	25 (28%)	6 (30%)	31 (28.44%)
11 to 20 years	24 (27%)	2 (10%)	26 (23.85%)
21 to 30 years	18 (20%)	3 (15%)	21 (19.27%)
31 to 40 years	7 (8%)	3 (15%)	10 (9.17%)
> 41 years	15 (17%)	6 (30%)	21 (19.27%)
Total	89 (100%)	20 (100%)	109 (100%)

Bacterial isolation peaked in the 0–20 age group, and as people age, the percentage of bacteria isolated falls. In the age groups of 0–10 and >40, the cultural negative was high.

**Table 6** Association of area and culture result

Area	Positive Culture: Number (%)	Negative culture: Number (%)	Total: Number (%)
Urban	33 (38.4%)	0 5 (21.7%)	38 (34%)
Rural	53 (61.6%)	18 (78.3%)	71 (66%)
Total	86 (100%)	23 (100%)	109 (100%)

The above table shows high culture positivity (61.6%) in rural cases than urban cases (38.4%).

Pattern of Organisms	Sex: Male/Female		Total Number (%)
	Male: Number (%)	Female: Number (%)	
<i>Staphylococcus aureus</i>	26 (36.1%)	10 (35.7%)	36 (36%)
<i>Coagulase negative staphylococci</i>	7 (9.7%)	2 (7.1%)	9 (9%)
<i>Streptococcus spp</i>	4 (5.6%)	0 (0.0%)	4 (4%)
<i>Pseudomonas aeruginosa</i>	26 (36.1%)	8 (28.6%)	34 (34%)
<i>Citrobacter freundii</i>	3 (4.2%)	1 (3.6%)	4 (4%)
<i>E.coli</i>	1 (1.4%)	3 (10.7%)	4 (4%)
<i>Klebsiella spp</i>	1 (1.4%)	1 (3.6%)	2 (2%)
<i>Proteus Mirabilis</i>	1 (1.4%)	1 (3.6%)	2 (2%)
<i>Acinetobacter spp.</i>	3 (4.2%)	1 (3.6%)	4 (4%)
<i>Enterobacter spp</i>	0 (0.0%)	1 (3.6%)	1 (1%)
Total	72 (100%)	28 (100%)	100 (100%)


According to the above table, *Staphylococcus auerus* 10 (37%) was the most common organism in females, while *Pseudomonas aueruginosa* and *Staphylococcus auerus* 26 (38%) each had equal incidence in males.

**Table 8** Area wise prevalence of organisms

Organisms isolated	Area: Rural/Urban		Total numbers(%)
	Rural: Number(%)	Urban: Number(%)	
<i>Staphylococcus aureus</i>	19 (30.2%)	17 (45.9%)	36 (36%)
<i>Coagulase negative staphylococci</i>	6 (9.5%)	3 (8.1%)	9 (9%)
<i>Streptococcus spp</i>	4 (6.3%)	0 (0.0%)	4 (4%)
<i>Pseudomonas aeruginosa</i>	22 (34.9%)	12 (32.4%)	34 (34%)
<i>Citrobacter freundii</i>	3 (4.8%)	1 (2.7%)	4 (4%)
<i>E.coli</i>	2 (3.2%)	2 (5.4%)	4 (4%)
<i>Klebsiella spp</i>	2 (3.2%)	0 (0.0%)	2 (2%)
<i>Proteus Mirabilis</i>	1 (1.6%)	1 (2.7%)	2 (2%)
<i>Enterobacter spp</i>	1 (1.6%)	0 (0.0%)	1 (1%)

<i>Acinetobacter spp.</i>	3 (4.8%)	1 (2.7%)	4 (4%)
Total	63 (100%)	37 (99.9%)	100 (100%)

*Staphylococcus aureus* 17 (40.5%) was the most common bacterium recovered from patients in rural areas, whereas *Pseudomonas aeruginosa* 21 (44.8%) was identified from patients in urban areas.



According to the above table, *Pseudomonas* species and *Staphylococcus aureus* were the most prevalent bacteria in

**Table 9** Age wise prevalence of organisms

Organisms isolated	Age Group					Total
	0-10	11-20	21-30	31-40	>40	
	Number (%)	Number (%)	Number (%)	Number (%)	Number (%)	Number (%)
<i>Staphylococcus aureus</i>	11 (39.3%)	9 (30.0%)	7.0 (41.2%)	3.0 (50.0%)	6.0 (31.6%)	36.0 (36%)
<i>Coagulase negative staphylococci</i>	3 (10.7%)	2 (6.7%)	1.0 (5.9%)	1.0 (16.7%)	2.0 (10.5%)	9.0 (9%)
<i>Streptococcus spp</i>	2 (7.1%)	2 (6.7%)	0.0 (0.0%)	0.0 (0.0%)	0.0 (0.0%)	4.0 (4%)
<i>Pseudomonas aeruginosa</i>	9 (32.1%)	15 (50.0%)	4.0 (23.5%)	1.0 (16.7%)	5.0 (26.3%)	34.0 (34%)
<i>Citrobacter freundii</i>	0 (0.0%)	0 (0.0%)	2.0 (11.8%)	1.0 (16.7%)	1.0 (5.3%)	4.0 (4%)
<i>E.coli</i>	2 (7.1%)	0 (0.0%)	1.0 (5.9%)	0.0 (0.0%)	1.0 (5.3%)	4.0 (4%)
<i>Proteus Mirabilis</i>	1 (3.6%)	0 (0.0%)	0.0 (0.0%)	0.0 (0.0%)	1.0 (5.3%)	2.0 (2%)
<i>Klebsiella spp</i>	0 (0.0%)	0 (0.0%)	0.0 (0.0%)	0.0 (0.0%)	2.0 (10.5%)	2.0 (2%)
<i>Enterobacter spp</i>	0 (0.0%)	0 (0.0%)	1.0 (5.9%)	0.0 (0.0%)	0.0 (0.0%)	1.0 (1%)
<i>Acinetobacter spp.</i>	0 (0.0%)	2 (6.7%)	1.0 (5.9%)	0.0 (0.0%)	1.0 (5.3%)	4.0 (4%)
Total	28 (99.9%)	30 (100%)	17.0 (100%)	6.0 (100%)	19.0 (100%)	100 (100%)



both adults and children. *Staphylococcus aureus* was the most prevalent pathogen in the adults.

Antibiotic tested	Sensitivity profile: Number of isolates sensitive to the tested antibiotic
Amoxyclav (1 µg)	21(21%)
Gentamicin(10 µg)	47(47%)
Cefotaxime(10µg)	46(46%)
Co trimaxazole(25 µg)	48(48%)
Amikacin(30 µg)	73(73%)
Ciprofloxacin(5 µg)	33(33%)

The above table shows the antibiotic susceptibility pattern of organisms isolated.

## DISCUSSION

The current investigation revealed that the highest proportion of patients fell within the age range of 1 to 10 years, accounting for 31 individuals, or 28.4% of the total cohort. Which is correlating with study by A S Adoga et al in a total of 80 patients 24(30%) children were below 10 years of age<sup>15</sup>. Mohammed Jamiu Kazeem, Ruqayyah Aiyaleso et al studied 380 patients with CSOM, maximum number patients were in the age group of 0-10(67.4%)<sup>10</sup>. A cross sectional study by Ashib Parvez et al, majority of the children were in the age group of 8 to 10 years(38.5%) followed by 6 to 8 years(36.4%)<sup>11</sup>. In a study by Y.K.Harshika et al, highest incidence in children with age group of 1 to 10 years (20.76%) followed by 11 to 20 years (18.4%) age group<sup>12</sup>.

Geeta S. H. *et al.* examined 250 CSOM samples; 30.4% of the patients in their study were between the ages of 0 and 10<sup>13</sup>. The current study and this study are related. In the current study, males were more impacted (64%) than females (36%). 204 CSOM samples were examined by Rajat Prakash *et al.* They found that women (53.92%) were more likely than men (46.08%) to be impacted<sup>14</sup>. Additionally, a research by A. S. Adoga et al. that included 30 males (37.5%) and 50 females (62.5%) found that females were more impacted than males<sup>15</sup>.

When Swarooparaniet al. examined 99 CSOM patients, they found that men were more likely to be impacted (56.6%) than women (43.4%)<sup>6</sup>. Of the 62 patients in a research by Sanjay Kumar *et al.*, 38 (61.29%) were men and 24 (38.7%) were women<sup>16</sup>. In their study of 120 CSOM patients, Vijay Kumar Poorey et al. found that 68 (56.66%) of them were men and 52 (43.34%) were women<sup>4</sup>. A study by Y.K. Harshika et al. found that men were more likely than women to have CSOM (53.07%)<sup>12</sup>. Additionally, S. Nikakhlagh *et al.*'s study revealed that men were more prevalent (54%) than women (46%)<sup>17</sup>. These studies support the current investigation.

According to the current study, the incidence of CSOM cases was higher in rural areas (66%) than in urban areas (34%). According to a study by Ashib Parvez *et al.*, 6.1% of the 630

children in the study had CSOM, and 7% of them lived in rural areas. and 1.8% came from cities. It was discovered to be connected with the child's nutritional status and the standard of life index<sup>11</sup>. Additionally, among the 125 patients in a research by Arti Agrawal et al., 70 (56.0%) came from rural areas and 55 (44.0%) from metropolitan areas<sup>18</sup>. Our study was associated with this one.

The most prevalent pathogens in both children and adults in the current investigation were *Staphylococcus aureus* and *Pseudomonas* species. This finding correlated with Geeta S H et al<sup>13</sup> and Swarooparaniet al<sup>6</sup>.

Unilateral infection was more prevalent in this study (82%) than bilateral infection (18%). Adoga et al., examined 80 CSOM patients and discovered bilateral CSOM in 6 cases and unilateral CSOM in 74 cases<sup>9</sup>. In a study of 380 patients, Mohammed Jamiu Kazeem, Ruqayyah Aiyaleso, et al. found that 3.4% of cases had unilateral CSOM and 96.6% had unilateral<sup>10</sup>. 120 of the 127 ear swabs in a research by Vijay Kumar Poorey et al. were bilateral, and seven of them were bilateral<sup>4</sup>. In 2009, Swarooparaniet al. conducted a study on 80 CSOM patients, reporting 76.8% unilateral and 23% bilateral<sup>6</sup>. Additionally, Sanjay Kumar et al. examined 62 individuals; of these, 54 (87.09%) had unilateral disease and 8 (12.09%) had bilateral disease<sup>16</sup>. These results are in line with current.

23 (21%) of the specimens in the current investigation tested negative for the culture, while 86 (79%) tested positive. The other studies likewise have different cultural outcomes. 116 (91.34%) of the 127 ear swabs that were taken for the study by Vijay Kumar Poorey et al. produced microbial growth. This finding may be explained by the efficacy of the previous antibiotic treatment, inadequate methods for collecting ear swabs, and sending swabs for cultures when the diagnosis is unclear<sup>4</sup>. In a study by Sateesh Kumar Malkappa *et al.*, 130 samples were included. of which 126 samples (96.92%) tested positive for culture<sup>8</sup>. 130 instances were examined by Y.K. Harshika et al. Of the 130 instances, 14 (10.7%) had no growth or skin contaminants, and 116 (89.5%) were culture positive<sup>12</sup>. 186 CSOM patients were investigated by Rajat Prakash et al. They found that 18 (8.82%)% of the cases in their study had negative cultures<sup>14</sup>. This study is consistent with ours.

Out of 86 culture-positive cases, 100 bacterial isolates were obtained. *Staphylococcus aureus* 36 (36%) was the most frequently isolated organism, followed by *Pseudomonas* spp. 34 (34%), and Coagulase-negative *Staphylococcus* 9 (9%). The remaining four percent each are *Streptococcus* spp., *Acinetobacter* spp., *Citrobacter* spp., and *E. coli*. *Enterobacter* spp. were identified from one sample (1%), whereas *Klebsiella* spp. and *Proteus vulgaris* 2 (2% each) were isolated.

*Pseudomonas* spp. was the most common gram-negative bacillus recovered in the investigation by Mohammed Jamiu Kazeem et al<sup>10</sup>. According to research by Kenna *et al.*<sup>19</sup>, Fliss et al.<sup>20</sup>, Papastavros *et al.*<sup>21</sup>, Arguedas as al<sup>22</sup> and Brook *et al.*<sup>23</sup>, *Pseudomonas* spp. was the most common bacteria recovered from the patients with CSOM.

32.4% of the *Staphylococcus aureus* in the S. Nikhakhlagh et al. investigation were followed by *Pseudomonas aeruginosa* (21.6%)<sup>17</sup>. *S. aureus* 47 (37.6%) was the most common organism in a research by Arti Agrawal *et al.*, followed by *P. aeruginosa* 41 (32.8%), *Klebsiella pneumoniae* 5 (4%), *E. coli* 4 (3.2%), *Streptococcus pneumoniae* 2 (1.6%), and *Proteus mirabilis* 1 (0.8%)<sup>18</sup>. Our research findings are consistent with those of the aforementioned studies. Vijay Kumar Poorey et al. examined 120 cases, 116 of which were culture-positive. They discovered that *P. aeruginosa* spp. (37.21%) was the most common bacterium, followed by *Staphylococcus* spp. (24.81%)<sup>4</sup>. *Proteus mirabilis* and *Citrobacter* spp. 8 (5.75%) each, *E. coli* (3.59%), *Enterobacter* spp. 3 (2.15%), *Acinetobacter* spp. CONS2 (1.43%) each, *Staphylococcus aureus* 30 (21.58%), *Klebsiella* spp. 13 (9.35%), and *P. aeruginosa* 46 (30.09%) were the most frequently isolated aerobic organisms in a study by Y K. Harshika *et al.*<sup>12</sup>.

## CONCLUSION

In our nation's rural areas, the prevalence of CSOM is still significant and typically found in younger age groups. Complication rates can be decreased by early detection of CSOM and proper treatment. Controlling infection presents the most treatment challenge because of its recurrent nature and the emergence of resistant pathogenic organisms. Before beginning empirical therapy, it is crucial to understand the local microbial flora in CSOM.

## Statements and Declarations

Funding: "This research received no external funding"

Conflicts of Interest: "The authors declare no conflict of interest."

Acknowledgments: nil

## Reference

1. R.K. Sanjana, Y.I. Singh, N.S. Reddy. Aerobic bacteriology of chronic suppurative otitis media (CSOM) in tertiary care hospital: A retrospective study. Journal of college of Medical sciences-Nepal. 2011;7: p.1-8.
2. Monasta L, Ronfani L, Marchetti F, Montico M, Vecchi Brumatti L, Bavcar A, et al. Burden of disease caused by otitis media: systematic review and global estimates. PloS One. 2012;7(4):e36226. <https://doi.org/10.1371/journal.pone.0036226>
3. World Health Organization. Chronic suppurative otitis media: burden of illness and management options. 2004: <https://apps.who.int/iris/handle/10665/42941>.
4. Vijay Kumar Poorey, Pooja Thakur. Clinicomicrobiological evaluation and antibiotic susceptibility in cases of chronic suppurative otitis media. Indian Journal of Otolaryngology. April 2015 ; Vol 21; Issue 2:107-110.
5. Monique Verhoeff, Erwin L. van der Veen, Maroeska M. Rovers, Elisabeth A.M. Sanders, Anne G.M. Schilder. Chronic suppurative otitis media: A review. International journal of pediatric otorhinolaryngology. 2006;: p.1-12.
6. Swaroopani NB, Kardesai SG and Metgud SC. Aerobic bacteriological study of Chronic suppurative otitis media with special reference to MRSA and ESBL. SMU Medical Journal. 2014 January; 1: p. 119-128.
7. Saurabh Varshney. Deafness in India. Indian Journal of Otolaryngology. April 2016 ; Vol 22; Issue 2:73-76.
8. Sateesh Kumar Malkappa, Saileela Kondapaneni et al. Study of aerobic bacteriological isolates and their antibiotic susceptibility pattern in chronic suppurative otitis media. Indian Journal of Otolaryngology. July 2012; Vol 18; Issue 3:136-139.
9. AS Adoga, EN Ma'an, D Malu et al. Swab and aspiration specimen collection methods and antibiogram in chronic suppurative otitis media at Jos University Teaching Hospital: Which is superior?. Annals of African Medicine. 2010; Volume 9; Issue 4; Page 230-234.
10. Mohammed Jamiu Kazeem, Ruqayyah Aiyaleso. Current bacteriological profile of chronic suppurative otitis media in a tertiary facility of Northern Nigeria. Indian Journal of Otolaryngology. July 2016 ; Vol 22 ; Issue 3: 157-161.
11. Ashib Parvez, Zulfia Khan, SF Hashmi et al. A cross sectional study of chronic suppurative otitis media and its associated factors among primary school children in rural and urban areas of Aligarh, India. Int J Community Med Public Health. 2016 Aug;3(8):2150-2154.
12. Y. K. Harshika, S. Sangeetha and R. Prakash. Microbiological Profile of CSOM and their Antibiotic Sensitivity Pattern in a Tertiary Care Hospital. Int.J.Curr.Microbiol. App.Sci (2015) 4(12): 735-743
13. S.H, Geeta. "Study of aerobes, Anaerobes & Fungi in CSOM in a referral Hospital of Bangalore Rural". Journal of Evaluation of medical and Dental Sciences. 2014 June 09; 3(23): p. 6297-6303.
14. Rajat Prakash, Deepak Juyal, Vikrant Negi, Shekhar Pal, Shamanthadekhandi, Munesh Sharma, and Neelam Sharma. Microbiology of chronic suppurative otitis media in tertiary care set up of Uttarakhand state, India. North American Journal of Medical Sciences. 2013 April;: p. 282-287.
15. AS Adoga, EN Ma'an, D Malu et al. Swab and aspiration specimen collection methods and antibiogram in chronic suppurative otitis media at Jos University Teaching Hospital: Which is superior?. Annals of African Medicine. 2010; Volume 9; Issue 4; Page 230-234.
16. Sanjay Kumar, Ravinder Sharma et al. Bacterial flora of infected unsafe CSOM. Indian Journal of Otolaryngology. October 2012; Vol 18; Issue 4:208-211.
17. S. Nikhakhlagh, A. D. Khoskravi, A. Fazlipour, M.

- Safarzadeh and N Rashidi. Microbiological findings in patients with chronic suppurative otitis media. *J.Med. Sci.* 2008 August 01; 8(5): 503-506.
18. Arti Agrawal, Dharmendra Kumar, Ankur Goyal et al. Microbiological profile and their antimicrobial sensitivity pattern in patients of otitis media with ear discharge. *Indian Journal of Otolaryngology*. January 2013; Vol 19; Issue 1:5-8.
  19. M.A. Kenna, C.D. Bluestone, J.S. Reilly, R.P. Lusk, Medical management of chronic suppurative otitis media without cholesteatoma in children, *Laryngoscope* 96 (2) (1986) 146-151.
  20. D.M. Fliss, R. Dagan, N. Meidan, A. Leiberman, Aerobic bacteriology of chronic suppurative otitis media without cholesteatoma in children, *Ann. Otol. Rhinol. Laryngol.* 101 (10) (1992) 866-869.
  21. T. Papastavros, H. Giamarellou, S. Varlejides, Role of aerobic and anaerobic microorganisms in chronic suppurative otitis media, *Laryngoscope* 96 (4) (1986) 438-442.
  22. A. Arguedas, C. Loaiza, J.F. Herrera, E. Mohs, Antimicrobial therapy for children with chronic suppurative otitis media without cholesteatoma, *Pediatr. Infect. Dis. J.* 13 (10) (1994) 878-882.
  23. I. Brook, Prevalence of beta-lactamase-producing bacteria in chronic suppurative otitis media, *Am. J. Dis. Child* 139(3) (1985) 280-283.
  24. M. de Uzeda, E.R. Rocha, Bacteriology of chronic otitis media affecting children living in Rio de Janeiro, *Ear Nose Throat J.* 68 (6) (1989), 448, 450, 452.
  25. N. Miro, Controlled multicenter study on chronic suppurative otitis media treated with topical applications of ciprofloxacin 0.2% solution in single-dose containers or combination of polymyxin B, Neomycin, and hydrocortisone suspension, *Otolaryngol. Head Neck Surg.* 123 (5) (2000) 617-623.
  26. Gerald B. Healy Otitis media and middle effusions. *Otorhinolaryngology, Head and Neck Surgery*. John Jacob Ballenger. 17th edition, Williams and Wilkins 2009. 1727, 217-226, 229-238
  27. Ananthanarayan R, Jayaram Paniker CK. Textbook of microbiology. 2009; 8th edition. 195-202, 270-282, 315-318, 667-668
  28. Mackie and McCartney's Practical medical microbiology. Collee JG, Duguid JP, Fraser AG, Marmion BP, editors. Edinburgh: Churchill Livingstone ELSEVIER; 2006. 14th edition. 131-149, 245-262, 413-422, 361-383.
  29. Clinical and Laboratory Standards Institute antimicrobial susceptibility testing standards. 2016, 26th ed. 137-143.
  30. Child and Adolescent Health and Development Prevention of Blindness and Deafness. World Health Organization Geneva, Switzerland. 2004. WHO Library Cataloguing-in-Publication Data Chronic suppurative otitis media: burden of illness and management options. 2004, 38-67.
  31. Bergey's Manual of determinative bacteriology. 9th edition. In: Holt TG, Krieg MR, Sneath PHA, Staley JT, Williams ST, editors. Baltimore: Lippincott Williams and Wilkins; 1994. 332-334, 364-368, 454-464.
  32. Lippincott Williams and Wilkins. Koneman's color atlas and textbook of diagnostic microbiology. 6th edition. 624-662.

**How to cite this article:**

Jyoti R. Hundekar., Surekha Y.A and S. Sadiqa Begum. (2025). Chronic suppurative otitis media infections: bacterial profile and demographic features among patients attending tertiary care hospital. *Int J Recent Sci Res.* 16(01), pp.050-056.

\*\*\*\*\*